

Characterization of Bioactives and Nutra-Pharmaceutical Potential of Supercritical Fluid and Hydro-Distilled Extracted Coriander Leaves Essential Oil

Dose-Response:
An International Journal
October-December 2022:1–10
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15593258221130749
journals.sagepub.com/home/dos


Ali Abbas^{1,2}, Farooq Anwar², Naveed Ahmad³ , Muhammad Shahid⁴, Samiah H. Al-Mijalli⁵, Muhammad Yaseen³, Sidra Farooq⁶, and Munawar Iqbal³ 

Abstract

The volatiles chemical composition and biological attributes of coriander (*Coriandrum sativum* L.) leaves essential oil obtained by two extraction techniques namely supercritical fluid extraction and hydro-distillation is appraised. The coriander essential oil yield (.12%) by hydro-distillation was slightly higher than that of supercritical fluid extraction (.09%). The physico-chemical variables of the essential oil obtained from both the techniques varied in significantly ($P < .05$). GC-MS analysis identified 23 different components in supercritical fluid extracted oil and 18 components in hydro-distilled essential oil having linalool as major component (51.32% and 61.78%, respectively) followed by phytol (12.71%). The oil recovered by supercritical fluid extraction exhibited greater DPPH radical scavenging activity as well as reducing power as compared to the essential oil obtained by hydro-distillation technique along with a stronger biofilm inhibition and least hemolysis. The results of antimicrobial activity revealed that super critical fluid extracted essential oil has potent antifungal and antibacterial activity against *P. multocida* and *A. alternata*, whereas hydro-distilled essential oil displayed better antimicrobial potential against *E. coli* and *A. niger*. Overall, these results depict that supercritical fluid extraction is superior than hydro-distillation with regard to isolation of better-quality coriander essential oil for nutra-pharmaceutical developments.

Keywords

hydro-distilled, coriander oil, supercritical fluid extraction, linalool, gas chromatography-mass spectrometry, natural antioxidant, antimicrobials agents

Introduction

Coriander (*Coriandrum sativum* L.), belonging to the family *Apiaceae* (Umbelliferae), has been described as one of the oldest herbs used for culinary and medicinal purposes.^{1,2} A native to Italy, this species is now broadly cultivated in different zones of the world such as North Africa, Russia, Central and Eastern Europe, Asia (Pakistan, India, Bangladesh, and China), and Mediterranean regions (Egypt, Morocco, and Malta).^{3,4} It can be grown throughout the year, but the best growth is observed between the months of October and February and flowers are produced between June and July.⁵ Because of its uses as a flavoring agent in foods and cosmetics, this annual herb has a greater economic value. The fruit and seeds of coriander are aromatic, having a mild bittersweet and spicy taste.⁶ The dried

¹ Department of Chemistry, Govt. Postgraduate Taleem-ul-Islam College, Chiniot, Pakistan

² Institute of Chemistry, University of Sargodha, Sargodha, Pakistan

³ Department of Chemistry, Division of Science and Technology, University of Education, Lahore, Pakistan

⁴ Department of Biotechnology and Bioinformatics, Government College University Faisalabad, Pakistan

⁵ Department of Biology, College of Sciences, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia

⁶ School Education Department, Pakistan

Corresponding Authors:

Munawar Iqbal, Department of Chemistry, Division of Science and Technology, University of Education, Lahore 54000, Pakistan.
Email: bosalvee@yahoo.com

Farooq Anwar, Institute of Chemistry, University of Sargodha, Sargodha, 41000, Pakistan.
Email: fqanwar@yahoo.com

Naveed Ahmad, Department of Chemistry, Division of Science and Technology, University of Education, Lahore 38000, Pakistan.
Email: mnavedahmad@yahoo.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE

and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

coriander seeds are employed as a component of curry powder and these are also used in stews and minced meat dishes. The fresh leaves of coriander have often been utilized in salads, making sauces and to garnish the food due to their characteristic aroma and attractive green color.^{7,8}

The main producers of coriander are Czech Republic, Poland, Romania, Mexico, Argentina, India, and Guatemala.⁵ Countries like Ukraine and India have the major control over the global market and trade of coriander oil.⁹ The oils obtained from coriander are also used in body care products, perfumes, cosmetics, and cosmeceuticals.¹⁰ The coriander seed and fresh leaves were studied to check the occurrence of different biologically active ingredients which impart various biological and health promoting attributes.^{2,11,12} Conventionally, different parts and products of coriander plant have been used to treat gastrointestinal disorder like dyspepsia, diarrhea, pain, anorexia, flatulence, and vomiting.²

A variety of medicinal herbs have been tested for essential oil (EO) production and commercialization.^{13,14} In particular, various EOs can be produced from different parts, that is, aerial (flower, buds, stem, leaves, fruits, peels, bark, wood, and seed) and underground (roots) of aromatic plants.^{15,16} The majority of odoriferous plants have been investigated to contain EO which can be obtained from dried, partially dehydrated or fresh parts of plants.¹⁶ Chemically, plants EOs mainly contain terpenoids (volatile constituents) which are of great biological importance due to their multiple pharmaceutical applications.¹⁶ The main applications of EOs include their uses as preservatives and flavoring agents, as well as potent bioactive agents.^{17,18} Besides the traditional use of EOs as health and healing agents in aromatherapy, these have also been documented as potential ingredient in preparation of flavored foods, perfumery products, cosme-nutraceuticals, and natural therapeutics.^{17,19}

The processing methods used for the handling and storage of plant materials as may affect the quality and quantity of EOs isolated.²⁰ The plant's genetic, environmental and physiological factors as well as extraction techniques employed can alter the volatiles composition and ultimately the yield and quality of EO extracted from different sources.²¹ Among different techniques, SCFE (supercritical fluid extraction) is employed for the extraction of EOs from different plant materials including basil,¹⁷ sweet gale,²² and jasmine.²³ In this green extraction method, the aromas from the plants are extracted/isolated using supercritical fluid such as liquid CO₂. Carbon dioxide is an inert gas having mild values of critical temperature (31.1°C) and critical pressure (7.38 MPa) and is considered to be non-hazardous and non-toxic. The supercritical fluid extraction method is more selective and safer as compared to conventional methods (hydro-distillation) and the extracted product contains no residual solvent but have value added benefits.²⁴

The Soon valley, situated in the Punjab province of Pakistan, with Sakesar as the highest peak in the Salt Range, covers around 300-square-mile area. The valley has impressive scenic beauty due to presence of several waterfalls, natural lakes and ponds, and lush green mountains. The agro-

ecological and agro-climatic characteristics of the Soon valley, in terms of mild temperature, distinct topography, and low rainfall, are unique as compared with the neighboring regions resulting in unique phytochemical composition of the medicinal flora in this specific area.²⁵

In one of our recent studies, we appraised and compared the volatiles composition and biological principles of basil (*Ocimum basilicum* L.) essential oil obtained by different techniques.¹⁷ As a part of our ongoing research activities, in the present research work, we planned to appraise the unexplored biochemical profile of coriander leaves harvested from unique Soon valley of Punjab, Pakistan. So, this research project is specifically framed to appraise the biological attributes and composition of essential oil obtained from coriander plants grown in this region. As per our best understanding, this report is the first one which compares the yield, volatile bioactives profiling and biological attributes of coriander leaves essential oil produced by two different techniques, that is, hydro-distillation and SCFE.

Experimental

Sample Collection

The plant parts (stem and leaves) of coriander (*Coriandrum sativum* L.) were obtained during spring season from Soon Valley, District Khushab, Punjab, Pakistan (Coordinates: 32°58'N 72°15'E). A taxonomist from Botany Department, University of Sargodha, Sargodha, Pakistan, further authenticated the specimen. The collected samples were washed and dried (moisture content 6.50%) at room temperature prior to extraction of EO.

Chemical and Reagents

Free radical of 2,2-diphenyl-1-picrylhydrazyl (DPPH), trichloro-acetic acid, ammonium thio-cyanate, ascorbic acid (vitamin C), aluminum chloride (AlCl₃), ferrous chloride, potassium ferri-cyanate, sodium nitrite, ferric chloride, butylatedhydroxytoluene (BHT) (99.0 %), butylatedhydroxyanisole (BHA), dimethyl sulfoxide, 3-(4,5-dimethylthiazol-2,5-diphenyl tetrazolium bromide) (MTT), penicillin/ streptomycin solution, and fluconazole were supplied by Sigma Chemical Corporation, USA. Sterile resazurin tablets for the purpose of sterilization were obtained from BDH Laboratories. All other chemicals such as chloroform, ammonium thiocyanate, ferrous chloride, methanol, and sodium carbonate (anhydrous) used in the research were acquired from Merck, Germany, unless stated otherwise. Analytical grade reagents and chemical were used for this research work.

Recovery of EO

The pulverized coriander leaves samples were processed for the extraction of EO using two different techniques for

comparative study: conventional hydro-distillation (HD) technique and modern supercritical fluid extraction (SCFE) technique.

Hydro-Distillation

EO was recovered using Clevenger-type apparatus. For this purpose, plant sample (3 kg) was hydro-distilled for 3 hours. The recovered oil was dehydrated over sodium sulfate (anhydrous). After filtration, recovered oil was preserved at -4°C until analyzed for different experiments.^{16,17}

Supercritical Fluid Extraction (SCFE)

A commercial SCFE instrument (Deven' supercritical private, Ltd., India) was used for the recovery of EO from coriander plant material. The plant sample (5 kg) was placed in the extractor and heated for 60 minutes at 45°C before the starting the process. Flow rate of supercritical CO_2 was adjusted as 10 mL/minute, under 100 bar pressure at 45°C . The extraction was done for 90 minutes under static condition and then for 30 minutes under dynamic conditions. The EO was collected at outlet and preserved at -4°C for experimental uses.²⁶

Analysis of Essential Oils

Yield and Physical Analyses. The yield (%) of the isolated EO was determined by weight of initially used plant. The physical analyses of the EOs include color, solubility (in 70% alcohol), density (25°C), and refractive index (25°C). These analyses were conducted following Abbas et al.¹⁷ Density and refractive index of EO were measured by densimeter (Anton Paar, model DMA 602, Austria), digital refractometer RX-7000a (Atago Co. Ltd., Japan), respectively.

Gas Chromatography (GC). The bioactives chemical profiling of the extracted coriander EO was studied using gas chromatography (PerkinElmer, Model 8700, USA) coupled with FID detector. Capillary column (HP-5MS, 30 m length, .25 mm diameter) was used in which thickness of stationary phase was .25 μm . The temperature of the column was initially adjusted at 80°C (for 3 min), followed by a gradual increase in temperature by 4°C per minute till 220°C , and finally maintained at 220°C for 10 minutes. The mobile phase used was an inert noble gas (He) which was allowed to flow at the rate of 1.5 mL per minute. The sample of EO (1.0 μL) was introduced in the injector (220°C) in a split mode, while detector temperature was maintained at 290°C .

GC-MS Analysis. The volatile bioactive components were further authenticated by using another advanced technique gas chromatography coupled with mass spectrometer (GCMS) (Agilent, model 6890N, MS-5975). For this analysis, the type of column and its temperate programming was the same as previously used in GC-FID method along with the same carrier gas with same

flow rate, but here the detection of the components was performed using mass spectrometer with electron ionization mode (70 eV). Identification of separated components was carried out by comparing their mass spectra with those of standard spectra published in NIST mass spectral library and finding out the retention indices (RI) relative to the normal alkane ($\text{C}_9\text{--C}_{24}$) and compared with the RI values of authentic compounds.^{17,27}

Antioxidant Potential

The antioxidant potential of the extracted coriander EO was estimated by performing different assays including reducing potential and free radical scavenging capacity.

DPPH[•] Scavenging Assay. A stable free radical, DPPH[•], was employed to evaluate antioxidant ability of recovered EOs.¹⁷ Various concentration of sample (.5–100 $\mu\text{g/mL}$) were mixed with 1.0 mL dilute solution of DPPH free radical (90 μmolar) and rectified methanol (95%) to make the total volume of mixture up to 4 mL. Positive control (butylated hydroxy toluene) was processed for comparing with the activity of the sample essential oil. The final solution was incubated and OD was recorded at 515 nm followed by construction of a plot between inhibition (%) and concentration to calculate IC_{50} value (concentration that scavenged 50% of the free radical).

Estimation of Reducing Power. The tested EOs were assessed for their reducing ability using following the method used by Abbas et al.¹⁷ Different amounts of EOs ranging from 2.5 mg to 10 mg were reacted with Na phosphate buffer (.2 M, pH = 6.6, 5.0 mL) and potassium ferricyanide (1.0%, 5 mL). After incubating at 50°C , this mixture was mixed with trichloroacetic acid (10%, 5.0 mL) followed by centrifuging for 10 minutes to separate upper layer which was diluted with water (5.0 mL). The resultant solution was added to iron (III) chloride (.1%, 1.0 mL) to record its optical density (OD) at 700 nm.

Antimicrobial Potential of EOs

Antimicrobial activity of tested EOs was evaluated against different pathogenic strains of bacteria (*S. aureus*, *P. multocida*, *E. coli*, and *B. subtilis*) and fungus (*A. flavus*, *A. niger*, *A. alternate*, and *G. lucidum*,) were used. Nutrient agar (NA) at temperature of 37°C was used as culture medium to grow bacteria while potato dextrose agar (PDA) at 30°C was used for fungal growth. The growth medium was purchased from Oxoid (Hampshire, UK). Streptomycin was used as standard drug for bacterial strains while fluconazol was selected as standard drug for fungus.

Inhibition zone of the tested EO was measured by the disc diffusion method²⁸ by considering 6 mm of sample-soaked filter paper discs and placed on the already incubated culture of bacteria and fungus. The corresponding standard drugs were named as positive control; however, filter paper disc without sample was named as negative control. The inhibition

zone was calculated after 24 hours of incubation for bacterial strains and 3 days of incubation for fungal strains.

Furthermore, to calculate the minimum inhibitory concentration of EO that inhibited the growth of the microorganism, micro dilution broth method previously used by Abbas et al.¹⁷ was selected. The MIC (minimum inhibitory concentration) value (mg/mL) gives a good knowledge regarding the antimicrobial activity of the tested EO, smaller the MIC value greater is the antimicrobial activity.

Biofilm Inhibition

The biofilm inhibition of EOs was checked against *E. coli* and *S. aureus* following a modified method reported by Regev-Shoshani et al.²⁹ The cell suspension of microorganism (100 μ L) was added into 96-well micro titre plate along with the EO sample. After incubation for 3 days at 37°C, the liquid was removed and 100 μ L of crystal violet solution (1%) was added. After staining for 30 minutes, dye was removed, and wells were washed. It was incubated again for 15 minutes after adding rectified ethanol (95%) and then the absorbance was noted at 570 nm. Positive control (rifamycin) was also processed under similar experimental conditions.

Hemolytic Activity

The assay of hemolytic activity was performed *in vitro* on human erythrocytes of O blood group reported by Malagoli.³⁰ Blood sample was obtained from volunteers, centrifuged for 5 minutes at 5000 rpm, and 2.0% suspension of erythrocyte was made in phosphate buffer saline. In this process, .85% of NaCl solution was added into various concentration of the EO solutions (50–500 μ g/mL), followed by the addition of already prepared erythrocyte suspension, which was kept at room temperature for 30 minute, upper layer of the mixture was collected to record the OD at 540 nm. Positive control (triton X-100) and negative control (phosphate buffer saline) were processed under similar experimental conditions. Hemolytic activity (%) was estimated by comparing with the absorbance value of positive control.³¹

Statistical Analysis

Three different samples of coriander plant leaves/EOs were used in each extraction technique to perform replicate experiments. To evaluate significant difference ($P < .05$), ANOVA (analysis of variance) was conducted by STATISTICA v5.5 (Computer software). The data were computed as mean $\pm$ standard deviation.³²

Results and Discussion

Yield and Physico-chemical Analysis

The results for percentage yield and physical parameters of the tested coriander EO are given in Table 1. The EO yield (w/w)

(.12%) by hydro-distillation is slightly higher than that of essential oil yield by SCFE technique (.09%), though this difference is not significant. This low yield of coriander essential oil in our research work is quite comparable to that of a previous study³⁷ in which the yield was noted to be .15%. Furthermore, the color of EO obtained by SCFE was light yellow while by hydro-distillation was just colorless. The test of solubility revealed that the oil obtained by hydro-distillation has slightly higher solubility (2.5 volume in 70% alcohol) than its counterpart (2.1 volume in 70% alcohol). Similarly, the oil extracted by SCFE has slightly higher density (.87 g/mL at 25°C) and refractive index (1.3800) than the oil obtained by hydro-distillation, which offered showed a density of .84 g/mL at 25°C and refractive index of 1.3300. It is clear from the results of physical analysis that the oils obtained by both the techniques are not significantly different from each other, except the oil obtained by SCFE is slightly concentrated. The minute difference in the physico-chemical parameters of the EOs in relation to the techniques employed can be connected with variation in the EO's composition.³³ In the previous studies, it is also appraised that the EO obtained through multiple methods have a little difference in the physico-chemical properties.^{34,35}

Composition of Coriander Essential Oil

The chemical compositional data of coriander EO produced by SCFE and HD and scrutinized by GC-MS is presented in Table 2. In the EO extracted by SCFE, a total 23 different components were identified including the major component linalool (51.34%) and others such as phytol (12.71%), α -pinene (9.91%), methyl linolenate (6.19%), geranyl acetate (4.23%), and camphor (3.45%) (Figure 1). However, in the hydro-distilled essential oil, only 18 components were identified including the principal component such as linalool (61.78%) followed by α -pinene (8.89%), camphor (7.16%), geranyl acetate (5.87%), and nerol (3.15%). The contents of the major components of the tested *C. sativum* EOs in the present work were quite comparable with those reported earlier.³⁶ Anwar et al.³⁷ found that coriander EO contained linalool (69.60%) as a major component with considerable amount of geranyl acetate (4.99%), γ -terpinene (4.17%), and α -pinene (1.63%). In another study, Singh et al.³ identified total of 52 components from coriander seed EO grown in India. The results revealed that the composition of major components in this study was quite comparable to our present compositional data. In a report by Zoubiri and Baaliouame,³⁸ a total of 17 constituents were recognized in the coriander seed EO from Algeria and linalool was found to be the principal component with the percentage of 73.1%. According to the reports of Bhuiyan et al.,³⁹ 53 components were identified from the EO of coriander seeds, harvested in Bangladesh. Again, linalool was found as the chief component followed by geranyl acetate and gamma-terpinene as the second and third larger compounds. Such variation in the chemicals profiling of *C. sativum* EO

Table 1. Yield (%) and physico-chemical parameters of coriander EOs.

Technique	Yield (w/w) (%)	Color	Solubility	Density g/mL (25°C)	Refractive index (25°C)
SCFE*	.09 ± .01 ^a	Very light yellow	2.1 part in alcohol (70%)	.87 ± .03 ^a	1.38 ± .04 ^a
Hydro-distillation	.12 ± .02 ^a	Just colorless	2.5 part in alcohol (70%)	.84 ± .02 ^a	1.33 ± .05 ^a

Note: Data is reported as mean ± SD, superscript (a) within the same column show significant ($P < .05$) differences of means between the two essential oil extraction methods.

*SCFE = Supercritical fluid extraction.

Table 2. Percentage composition comparison of some major component in coriander EOs produced by SCFE and HD techniques.

Sr. #	Compound name	IUPAC name	RI*	SCFE**	Hydro-distillation	Method of identification
1	α -Pinene	2,6,6-trimethylbicyclo[3.1.1]hept-2-ene	941	9.91 ± .62 ^a	8.89 ± .45 ^b	a, b, c
2	Camphene	2,2-dimethyl-3-methylidenebicyclo[2.2.1]heptane	950	—	1.42 ± .07	a, b, c
3	α -Cymene	1-methyl-2-(propan-2-yl)benzene	1025	—	2.12 ± .09	a, b, c
4	Limonene	1-methyl-4-(1-methylethenyl)-cyclohexene	1035	1.43 ± .12 ^b	2.49 ± .12 ^a	a, b, c
5	γ -Terpinen	1-methyl-4-propan-2-ylcyclohexa-1,4-diene	1058	—	3.95 ± .21	a, b, c
6	Linalool	3,7-dimethylocta-1,6-dien-3-ol	1096	51.34 ± 1.91 ^b	61.78 ± 3.21 ^a	a, b, c
7	Camphor	1,7,7-trimethylbicyclo[2.2.1]heptan-2-one	1147	3.45 ± .22 ^b	7.16 ± .34 ^a	a, b, c
8	Nerol	(2Z)-3,7-dimethylocta-2,6-dien-1-ol	1325	—	3.15 ± .19	a, b, c
9	Geranyl acetate	[(2E)-3,7-dimethylocta-2,6-dienyl] acetate	1382	4.23 ± .31 ^b	5.87 ± .31 ^a	a, b, c
10	Methyl Linolenate	methyl (9Z,12Z,15Z)-octadeca-9,12,15-trienoate	2040	6.19 ± .62	—	a, b, c
11	Linoleic acid	(9Z,12Z)-octadeca-9,12-dienoic acid	2104	1.42 ± .21	—	a, b, c
12	Phytol	(E,7R,11 R)-3,7,11,15-tetramethylhexadec-2-en-1-ol	2128	12.71 ± .85	—	a, b, c
Total Components Identified				23	18	

Note: Data is reported as mean ± SD, superscript within the same row displays a significant ($P < .05$) differences of means between SCFE and HD.

*RI = Retention index; **SCFE = Supercritical fluid extraction.

^a = Identification by retention index.

^b = identification by comparing with authentic compounds.

^c = identification by comparing of mass spectra.

may be mainly caused due to different agro-climatic conditions of the harvesting regions which determine and shape the morphological features of the crop.

Antioxidant Potential

Antioxidant potential of the tested coriander EOs obtained by both the techniques was appraised by reducing capacity and free radical scavenging ability (Table 3). The results showed significant difference in the DPPH radical scavenging activity of the EO for both the techniques. The DPPH radical scavenging activity was presented in terms of IC₅₀ value in µg/mL, which showed that the SCF extracted EO has good DPPH radical scavenging activity with least IC₅₀ value (9.11 µg/mL) as compared to hydro-distilled essential oil, which gave IC₅₀ of 10.34 µg/mL. Interestingly, SCF extracted EO gave even better result as compared to the standard compound (BHT) used in the experiment.

In reducing power assay, the yellow color of ferric ions is changed into bluish green color of ferrous ions after reduction. The reducing potential of EO depends upon the color intensity of the final reaction mixture, which shows that there is a direct relationship between the color intensity and the antioxidant powder.¹⁷ A gradual increment in the absorbance was recorded with increasing concentration of sample. At the concentration of 10.0 mg/mL, SCF-extracted EO showed better reducing potential (1.32) as compared to hydro-distilled EO (1.20) but less than standard synthetic antioxidant compound, which showed the absorbance of 1.91 at concentration of 10.0 mg/mL.

Foudah et al.⁴⁰ reported in his research work that coriander essential oil is a promising reservoir of natural antioxidants. In another study,⁴¹ the antioxidant potential was determined in oil and extracts from seeds and leaves of coriander. Overall, from the results of antioxidant activity it can be concluded that EO obtained from SCFE has better antioxidant and nutra-




Figure 1. Structure of some major components of coriander essential oil

Table 3. Antioxidant potential of coriander EOs.

Extraction Technique	DPPH* Radical Scavenging Assay		Reducing Power (Absorbance values)			
			Concentration (mg/mL)			
	Inhibition (%)	IC ₅₀ , µg/mL	2.5	5.0	7.5	10.0
SCFE**	42 ± 2 ^b	9.11 ± .21 ^b	.57 ± .04 ^b	.83 ± .07 ^a	1.01 ± .09 ^a	1.32 ± .12 ^b
Hydro-distillation	49 ± 2 ^a	10.34 ± .32 ^a	.68 ± .08 ^a	.79 ± .04 ^a	.98 ± .08 ^a	1.20 ± .14 ^b
BHT***	—	10.21 ± .29 ^a	—	—	—	1.91 ± .21 ^a

Note: Data is reported as mean ± SD, superscript (a, b) within the same column show significant ($P < .05$) difference in values for SCFE and HD.

*DPPH = 2,2-diphenyl-1-picrylhydrazyl. **SCFE = Supercritical fluid extraction. ***BHT = Butylated hydroxy toluene.

Table 4. Antimicrobial potential of coriander EOs.

Microorganism	Inhibition zone (mm)			MIC (µg/mL)		
	SCFE*	Hydro distillation	Standard Drug**	SCFE	Hydro distillation	Standard Drug**
Bacteria						
<i>S. aureus</i>	9.00 ± .25 ^c	16.00 ± .42 ^b	28.00 ± 1.12 ^a	132 ± 5 ^a	129 ± 5 ^a	62 ± 2 ^b
<i>B. subtilis</i>	9.00 ± .21 ^c	17.00 ± .37 ^b	31.00 ± 1.18 ^a	126 ± 4 ^a	103 ± 3 ^b	73 ± 3 ^c
<i>P. multocida</i>	12.00 ± .27 ^c	18.00 ± .51 ^b	30.00 ± 1.12 ^a	115 ± 3 ^a	86 ± 2 ^b	54 ± 2 ^c
<i>E. coli</i>	11.00 ± .22 ^c	21.00 ± .53 ^b	29.00 ± 1.21 ^a	127 ± 4 ^a	72 ± 2 ^b	63 ± 3 ^c
Fungus						
<i>G. lucidum</i>	15.00 ± .52 ^c	17.00 ± .61 ^b	26.00 ± 1.08 ^a	93 ± 2 ^a	82 ± 2 ^b	73 ± 2 ^c
<i>A. flavus</i>	18.00 ± .68 ^b	15.00 ± .58 ^c	28.00 ± 1.15 ^a	86 ± 3 ^b	102 ± 4 ^a	78 ± 2 ^c
<i>A. niger</i>	14.00 ± .47 ^c	19.00 ± .57 ^b	30.00 ± 1.23 ^a	82 ± 2 ^a	74 ± 2 ^b	53 ± 2 ^c
<i>A. alternata</i>	20.00 ± .72 ^b	17.00 ± .67 ^c	24.00 ± 1.02 ^a	82 ± 2 ^b	92 ± 3 ^a	83 ± 3 ^b

Note: Data is reported as mean ± SD, superscript (a, b, c) within the same row show significant ($P < .05$) differences of means between the two essential oil extraction methods. ** Standard drug *Streptomycin* for bacteria while *Fluconazole* for fungus.

*SCFE = Supercritical fluid extraction.

Table 5. Bioactivities of coriander EOs.

Bioactivity	Microbe	Coriander EO		Rifamacin	Triton-x-100
		SCFE*	Hydro-distillation		
Biofilm Inhibition %	<i>S. aureus</i>	78.61 ± 2.62 ^b	52.92 ± 1.82 ^c	87.43 ± 3.89 ^a	—
	<i>E. coli</i>	37.96 ± 1.26 ^c	64.75 ± 1.92 ^b	88.92 ± 3.45 ^a	—
Hemolytic Assay %		4.86 ± .19 ^b	31.58 ± 1.92 ^a	—	100

Note: Data is reported as mean ± SD, superscript (a, b, c) within the same row show significant ($P < .05$) differences of means between the two essential oil extraction methods.

*SCFE = Supercritical fluid extraction.

pharmaceutical potential as compared to EO obtained by hydro-distillation, which reveals the superiority of SCFE techniques over traditionally used hydro-distillation technique.

Antimicrobial Activity

Antimicrobial activity of coriander EO was determined and the related results are presented in Table 4. The coriander EO extracted by SCFE showed maximum antibacterial activity (12 mm inhibition zone) against *P. multocida* with MIC (115 µg/mL) while the hydro-distilled EO had potent activity (21 mm inhibition zone) against *E. coli* with MIC = 72 µg/mL. On the other hand, SCF extracted EO also showed stronger antifungal activity (20 mm inhibition zone) against *A. alternata* with MIC = 82 µg/mL while the hydro-distilled EO was more resistant against *A. niger* with inhibition zone and MIC values of 19 mm and 74 µg/mL, respectively. However, in comparison with standard drugs, antimicrobial potency of EOs extracted either by SCFE or hydro-distillation possessed less antimicrobial activity than the standard compounds such as streptomycin and fluconazole that were used as antibacterial and antifungal drugs in the present experiments, respectively. Interestingly, in contrast to the observations of Cantore et al,⁴² the tested EOs were more resistant to gram negative bacteria such as *E. coli* than Gram positive bacteria. Such types of trends in antimicrobial potential might be ascribed to chemical composition, agro-ecological, agro-climatic conditions, and analytical control of the experimental assays.

Several essential oils, which are mainly composed of terpenoids, were also studied for the occurrence of phenolic antimicrobial components.^{15,17,43-45} The phenolic compounds present in the plant essential oils show higher antimicrobial effect as compared to aldehyde, alcohol, and ketones.^{46,47}

Other Bioactivities (Biofilm Inhibition and Hemolytic Activity)

To further elucidate biological attributes of tested coriander EOs, some other biological activities like biofilm inhibition and hemolytic activity was also measured. Biofilm can preserve the bacteria by hindering the effects of antibiotics and antioxidants and host immune responses.⁴⁸ Due to the

presence of bacterial biofilm and its relative permeability some antibiotics and antioxidants would become ineffective,⁴⁹⁻⁵⁴ so a good antioxidant agent must also possess biofilm inhibition activity. Data regarding the biofilm inhibition is given in Table 5, which shows that the maximum biofilm inhibition was given by SCF extracted EO against *S. aureus* (78.61%) but less than Rifamacin (87.43%), standard drug used in the experiment. The result of our research work is in line with those of Duarte et al,⁵⁵ who described that coriander EO displayed significant anti-biofilm potential.

It is observed that some antioxidant compounds may have good antioxidant activity but also possess hemolytic effect, due to this reason they may not be able to use in pharmaceutical preparation. Hence for a good and safer antioxidant, it must have least hemolytic attributes. Therefore, in this project, hemolytic activity of tested EO was assessed (Table 5). The results clearly showed that the EO obtained by SCFE method has far less hemolytic activity (4.86%) as compared to the EO extracted by hydro-distillation (31.58%). These findings also support that the SCFE technique is better than conventional hydro-distillation technique toward isolation of nutra-pharmaceutical grade EOs. There are no previous literature reports available with which we can compare our results of the hemolytic activity. Hence, the present investigation revealed that the coriander EO is safer since no toxicity was observed, which have potential application for the development of nutra-pharmaceutical agents. The plants produced bioactive agents and these bioactive agents can be used for different fields for practical applications.⁵⁶⁻⁵⁸

Conclusion

Although the yield and physico-chemical attributes of the coriander EOs tested were significantly varied for SCFE and HD techniques. However, the results revealed that the coriander EO obtained from both the techniques have significant variations with regard to the biological activities. The EO obtained by SCFE technique possesses better biological potential due to presence of higher quantity of different potent bioactive components in comparison to EO recovered by hydro-distillation. Thus, in terms of their nutra-pharmaceutical applications, SCFE can be considered to be superior extraction techniques for the recovery of biologically

active volatile components. These research findings might motivate more researchers to explore the essential oil and phytochemicals profiling and biological potential of other medicinal plants from Soon Valley, especially meeting the demand of local modern food and pharmaceutical industry. Such data might also encourage the local growers and community to cultivate and harvest such medicinal herbs on large scale and install on farm SCFE plants for isolation of valuable essential oils with multiple food and nutra-pharmaceutical applications.

Acknowledgments

The authors express their gratitude to Princess Nourah bint Abdulrahman University Researchers Supporting Project (Grant No. PNURSP2022R158), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2022R158) Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

ORCID iDs

Naveed Ahmad  <https://orcid.org/0000-0002-5847-903X>

Munawar Iqbal  <https://orcid.org/0000-0001-7393-8065>

References

1. Weisany W, Tahir NAR, Schenk PM. Coriander/soybean intercropping and mycorrhizae application lead to overyielding and changes in essential oil profiles. *Eur J Agron*. 2021;126: 126283. DOI: [10.1016/j.eja.2021.126283](https://doi.org/10.1016/j.eja.2021.126283).
2. Sahib NG, Anwar F, HGilani A, Hamid AA, Saari N, Alkharfy KM. Coriander (*Coriandrum sativum* L.): A potential source of high-value components for functional foods and nutraceuticals. A review. *Phytother. Res*. 2013;27(10):1439-1456. DOI: [10.1002/ptr.4897](https://doi.org/10.1002/ptr.4897).
3. Singh G, Maurya S, De Lampasona MP, Catalan CA. Studies on essential oils, Part 41. Chemical composition, antifungal, antioxidant and sprout suppressant activities of coriander (*Coriandrum sativum*) essential oil and its oleoresin. *Flavour Fragrance J*. 2006;21(3):472-479. DOI: [10.1002/ffj.1608](https://doi.org/10.1002/ffj.1608).
4. Sriti J, Wannes WA, Talou T, Mhamdi B, Cerny M, Marzouk B. Lipid profiles of Tunisian coriander (*Coriandrum sativum*) seed. *JAOCs (J Am Oil Chem Soc)*. 2010;87(4):395-400. DOI: [10.1007/s11746-009-1505-1](https://doi.org/10.1007/s11746-009-1505-1).
5. Peter KV, Babu KN. Introduction to herbs and spices: medicinal uses and sustainable production. In: *Handbook of herbs and spices*. Woodhead Publishing; 2012:1-16. DOI: [10.1533/9780857095688.1](https://doi.org/10.1533/9780857095688.1).
6. Ceska O, Chaudhary SK, Warrington P, Ashwood-Smith MJ, Bushnell GW, Poultont GA. Coriandrin, a novel highly photoactive compound isolated from *Coriandrum sativum*. *Phytochemistry (Oxf)*. 1988;27(7):2083-2087. DOI: [10.1016/0031-9422\(88\)80101-8](https://doi.org/10.1016/0031-9422(88)80101-8).
7. Pellegrini M, Rossi C, Palmieri S, et al. Salmonella enterica control in stick carrots through incorporation of coriander seeds essential oil in sustainable washing treatments. *Front Sustain Food Syst*. 2020;4:14. DOI: [10.3389/fsufs.2020.00014](https://doi.org/10.3389/fsufs.2020.00014).
8. Kamat A, Pingulkar K, Bhushan B, Gholap A, Thomas P. Potential application of low dose gamma irradiation to improve the microbiological safety of fresh coriander leaves. *Food Control*. 2003;14(8):529-537. DOI: [10.1016/S0956-7135\(02\)00116-0](https://doi.org/10.1016/S0956-7135(02)00116-0).
9. Burdock GA, Carabin IG. Safety assessment of coriander (*Coriandrum sativum* L.) essential oil as a food ingredient. *Food Chem Toxicol*. 2009;47(1):22-34. DOI: [10.1016/j.fct.2008.11.006](https://doi.org/10.1016/j.fct.2008.11.006).
10. Prachayasittikul V, Prachayasittikul S, Ruchirawat S, Prachayasittikul V. Coriander (*Coriandrum sativum*): A promising functional food toward the well-being. *Food Res Int*. 2018;105:305-323. DOI: [10.1016/j.foodres.2017.11.019](https://doi.org/10.1016/j.foodres.2017.11.019).
11. Kačaniová M, Galovičová L, Ivanišová E, Vukovic NL, Štefániková J, Valková V, et al. Antioxidant, antimicrobial and antibiofilm activity of coriander (*Coriandrum sativum* L.) essential oil for its application in foods. *Foods*. 2020;9(3):282. DOI: [10.3390/foods9030282](https://doi.org/10.3390/foods9030282).
12. Begnami AF, Duarte MCT, Furletti V, Rehder VLG. Antimicrobial potential of *Coriandrum sativum* L. against different *Candida* species in vitro. *Food Chem*. 2010;118(1):74-77. DOI: [10.1016/j.foodchem.2009.04.089](https://doi.org/10.1016/j.foodchem.2009.04.089).
13. Maksimovic Z, Stojanović D, Šošarić I, Dajić Z, Ristić M. Composition and radical-scavenging activity of *Thymus glabrescens* Willd (Lamiaceae) essential oil. *J Sci Food Agric*. 2008; 88(11):2036-2041. DOI: [10.1002/jsfa.3311](https://doi.org/10.1002/jsfa.3311).
14. Busatta C, Vidal RS, Popielski AS, et al. Application of *Origanum majorana* L. essential oil as an antimicrobial agent in sausage. *Food Microbiol*. 2008;25(1):207-211. DOI: [10.1016/j.fm.2007.07.003](https://doi.org/10.1016/j.fm.2007.07.003).
15. Anwar F, Ali M, Hussain AI, Shahid M. Antioxidant and antimicrobial activities of EO and extracts of fennel (*Foeniculum vulgare* Mill.) seeds from Pakistan. *Flavour Fragrance J*. 2009; 24(4):170-176. DOI: [10.1002/ffj.1929](https://doi.org/10.1002/ffj.1929).
16. Hussain AI, Anwar F, Sherazi STH, Przybylski R. Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Food Chem*. 2008;108(3):986-995. DOI: [10.1016/j.foodchem.2007.12.010](https://doi.org/10.1016/j.foodchem.2007.12.010).
17. Abbas A, Anwar F, Ahmad N. Variation in physico-chemical composition and biological attributes of common basil essential oils produced by hydro-distillation and super critical fluid extraction. *J. Essent. Oil-Bear. Plants*. 2017;20(1):95-109. DOI: [10.1080/0972060X.2017.1280418](https://doi.org/10.1080/0972060X.2017.1280418).

18. Hussain AI, Anwar F, Chatha SA, et al. Chemical composition and bioactivity studies of the essential oils from two *Thymus* species from the Pakistani flora. *LWT-Food Sci Technol*. 2013; 50(1):185-192. DOI: [10.1016/j.lwt.2012.06.003](https://doi.org/10.1016/j.lwt.2012.06.003).
19. Kapustová M, Granata G, Napoli E, et al. Nanoencapsulated Essential Oils with Enhanced Antifungal Activity for Potential Application on Agri-Food, Material and Environmental Fields. *Antibiotics*. 2021;10(1):31. DOI: [10.3390/antibiotics10010031](https://doi.org/10.3390/antibiotics10010031).
20. Van Vuuren SF, Viljoen AM, Özék T, Demirci B, Başer KHC. Seasonal and geographical variation of *Heteropyxis natalensis* essential oil and the effect thereof on the antimicrobial activity. *South Afr J Bot*. 2007;73(3):441-448. DOI: [10.1016/j.sajb.2007.03.010](https://doi.org/10.1016/j.sajb.2007.03.010).
21. Masotti V, Juteau F, Bessière JM, Viano J. Seasonal and phenological variations of the essential oil from the narrow endemic species *Artemisia molinieri* and its biological activities. *J Agric Food Chem*. 2003;51(24):7115-7121. DOI: [10.1021/jf034621y](https://doi.org/10.1021/jf034621y).
22. Sokolova M, Orav A, Koel M, Kailas T, Müürisepp M. Composition of the oil and supercritical fluid CO₂ extract of sweet gale (*Myrica gale* L.) fruits. *J Essent Oil Res*. 2005;17(2): 188-191. DOI: [10.1080/10412905.2005.9698871](https://doi.org/10.1080/10412905.2005.9698871).
23. Younis A, Mehdi A, Riaz A. Supercritical carbon dioxide extraction and gas chromatography analysis of *Jasminum sambac* essential oil. *Pakistan J Bot*. 2011;43:163-168. [http://pakbs.org/pjbot/PDFs/43\(SI\)/24.pdf](http://pakbs.org/pjbot/PDFs/43(SI)/24.pdf)
24. Wood JA, Bernards MA, Wan WK, Charpentier PA. Extraction of ginsenosides from North American ginseng using modified supercritical carbon dioxide. *J Supercrit Fluids*. 2006;39(1): 40-47. DOI: [10.1016/j.supflu.2006.01.016](https://doi.org/10.1016/j.supflu.2006.01.016).
25. Ahmad N, Zuo Y, Lu X, Anwar F, Hameed S. Characterization of free and conjugated phenolic compounds in fruits of selected wild plants. *Food Chem*. 2016;190:80-89. DOI: [10.1016/j.foodchem.2015.05.077](https://doi.org/10.1016/j.foodchem.2015.05.077).
26. Rezazadeh S, Baha-Aldini BZBF, Vatanara A, et al. Comparison of super critical fluid extraction and hydrodistillation methods on lavender's essential oil composition and yield. *J. Med. Plants*. 2008;7(25):63-68. <http://dorl.net/dor/20.1001.1.2717204.2008.7.25.23.9>
27. Graikou KVK, Ngassapa O, Runyoro D, Chinou I. Composition and antimicrobial activity of the essential oils of three *Satureja* species growing in Tanzania. *Food Chem*. 2007;103(2): 319-324. DOI: [10.1016/j.foodchem.2006.07.051](https://doi.org/10.1016/j.foodchem.2006.07.051).
28. Wayne PA. *NCCLS: National Committee for Clinical Laboratory Standards. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically*; 1999.
29. Regev-Shoshani G, Ko M, Miller C, Av-Gay Y. Slow release of nitric oxide from charged catheters and its effect on biofilm formation by *Escherichia coli*. *Antimicrob. Agents Chemother*. 2010;54(1):273-279. DOI: [10.1128/AAC.00511-09](https://doi.org/10.1128/AAC.00511-09).
30. Malagoli D. A full-length protocol to test hemolytic activity of palytoxin on human erythrocytes. *Invertebr Surviv J*. 2007;4(2): 92-94. <http://hdl.handle.net/11380/421488>
31. Gould LA, Lansley AB, Brown MB, Forbes B, Martin GP. Mitigation of surfactant erythrocyte toxicity by egg phosphatidylcholine. *J Pharm Pharmacol*. 2000;52(10):1203-1209. DOI: [10.1211/0022357001777333](https://doi.org/10.1211/0022357001777333).
32. Steel RG, Torrie JH. *Principles and Procedures of Statistics*. New York, NY: McGraw-Hill Book Co. Inc.; 1980:481.
33. Vijayalakshmi A, Tripathi R, Ravichandiran V. Characterization and evaluation of antidermatophytic activity of the essential oil from *Artemisia nilagirica* leaves growing wild in nilgiris. *Int J Pharm Pharmaceut Sci*. 2010;2:93-97. <https://innovareacademics.in/journal/ijpps/Vol2Suppl4/803.pdf>
34. Hassanein HD, El-Gendy AENG, Saleh IA, Hendawy SF, Elmissiry MM, Omer EA. Profiling of essential oil chemical composition of some Lamiaceae species extracted using conventional and microwave-assisted hydrodistillation extraction methods via chemometrics tools. *Flavour Fragrance J*. 2020; 35(3):329-340. DOI: [10.1002/ffj.3566](https://doi.org/10.1002/ffj.3566).
35. Turek C, Stintzing FC. Stability of essential oils: a review. *Compr Rev Food Sci Food Saf*. 2013;12(1):40-53. DOI: [10.1111/1541-4337.12006](https://doi.org/10.1111/1541-4337.12006).
36. Wahba HE, Abd Rabbu HS, Ibrahim ME. Evaluation of essential oil isolated from dry coriander seeds and recycling of the plant waste under different storage conditions. *Bull Natl Res Cent*. 2020;44(1):1-7. DOI: [10.1186/s42269-020-00448-z](https://doi.org/10.1186/s42269-020-00448-z).
37. Anwar F, Sulman M, Hussain AI, Saari N, Iqbal S, Rashid U. Physicochemical composition of hydro-distilled essential oil from coriander (*Coriandrum sativum* L.) seeds cultivated in Pakistan. *J Med Plants Res*. 2011;5(15):3537-3544. DOI: [10.5897/JMPR.9000978](https://doi.org/10.5897/JMPR.9000978).
38. Zoubiri S, Baaliouamer A. Essential oil composition of *Coriandrum sativum* seed cultivated in Algeria as food grains protectant. *Food Chem*. 2010;122(4):1226-1228. DOI: [10.1016/j.foodchem.2010.03.119](https://doi.org/10.1016/j.foodchem.2010.03.119).
39. Bhuiyan MNI, Begum J, Sultana M. Chemical composition of leaf and seed essential oil of *Coriandrum sativum* L. from Bangladesh. *Bangladesh J Pharmacol*. 2009;4(2):150-153. doi: [10.3329/bjpv.v4i2.2800](https://doi.org/10.3329/bjpv.v4i2.2800).
40. Foudah AI, Alqarni MH, Alam A, Salkini MA, Ahmed EO, Yusufoglu HS. Evaluation of the composition and in vitro antimicrobial, antioxidant, and anti-inflammatory activities of Cilantro (*Coriandrum sativum* L. leaves) cultivated in Saudi Arabia (Al-Kharj). *Saudi J Biol Sci*. 2021;28(6):3461-3468. DOI: [10.1016/j.sjbs.2021.03.011](https://doi.org/10.1016/j.sjbs.2021.03.011).
41. Wangenstein H, Samuelsen AB, Malterud KE. Antioxidant activity in extracts from coriander. *Food Chem*. 2004;88(2): 293-297. DOI: [10.1016/j.foodchem.2004.01.047](https://doi.org/10.1016/j.foodchem.2004.01.047).
42. Lo Cantore P, Iacobellis NS, De Marco A, Capasso F, Senatore F. Antibacterial activity of *Coriandrum sativum* L. and *Foeniculum vulgare* Miller var. *vulgare* (Miller) essential oils. *J Agric Food Chem*. 2004;52(26):7862-7866. DOI: [10.1021/jf0493122](https://doi.org/10.1021/jf0493122).
43. Abebe A, Abebe M, Mekonnen A. Assessment of antioxidant and antibacterial activities of crude extracts of *Verbena officinalis* linn root or atuch (Amharic). *Chem Int*. 2017;3:172-184.
44. Rauf A, Hassan SM, Rafique S. Nutra-pharmaceutical efficacy appraisal of *Lavandula stoechas* leaves extracts in different solvents. *Chem Int*. 2021;8(4):242-248.
45. Friday C, Akwada U, Igwe OU. Phytochemical screening and antimicrobial studies of *Azelia africana* and *Detarium microcarpum* seeds. *Chem Int*. 2018;4(3):170-176.

46. Tao R, Sedman J, Ismail A. Antimicrobial activity of various essential oils and their application in active packaging of frozen vegetable products. *Food Chem.* 2021;360:129956. DOI: [10.1016/j.foodchem.2021.129956](https://doi.org/10.1016/j.foodchem.2021.129956).
47. Tepe B, Daferera D, Sokmen A, Sokmen M, Polissiou M. Antimicrobial and antioxidant activities of the essential oil and various extracts of *Salvia tomentosa* Miller (*Lamiaceae*). *Food Chem.* 2005;90(3):333-340. DOI: [10.1016/j.foodchem.2003.09.013](https://doi.org/10.1016/j.foodchem.2003.09.013).
48. Darouiche RO, Raad II, Heard SO, et al. A comparison of two antimicrobial-impregnated central venous catheters. *NEJM.* 1999;340(1):1-8. doi:[10.1056/NEJM199901073400101](https://doi.org/10.1056/NEJM199901073400101).
49. Nuță DC, Limban C, Chiriță C, et al. Contribution of Essential Oils to the Fight against Microbial Biofilms—A Review. *Processes.* 2021;9(3):537. DOI: [10.3390/pr9030537](https://doi.org/10.3390/pr9030537).
50. Dolatabadi M, Świergosz T, Ahmadzadeh S. Electro-Fenton approach in oxidative degradation of dimethyl phthalate-The treatment of aqueous leachate from landfills. *Sci Total Environ.* 2021;772:145323.
51. Dolatabadi M, Ghaneian MT, Wang C, Ahmadzadeh S. Electro-Fenton approach for highly efficient degradation of the herbicide 2, 4-dichlorophenoxyacetic acid from agricultural wastewater: Process optimization, kinetic and mechanism. *J. Mol. Liquids.* 2021;334:116116.
52. Ahmadzadeh S, Kassim A, Abdollahi YA. Conductometric Study of Complexation Reaction Between Meso-octamethylcalix [4] pyrrole with Titanium Cation in. *Int J Electrochem Sci.* 2011;6:4749-4759.
53. Ahmadzadeh S, Rezayi M, Karimi-Maleh H, Alias Y. Conductometric measurements of complexation study between 4-Isopropylcalix [4] arene and Cr³⁺ cation in THF–DMSO binary solvents. *Measurement.* 2015;70:214-224.
54. Rezayi M, Heng LY, Kassim A, Ahmadzadeh S, Abdollahi Y, Jahangirian H. Immobilization of tris (2 pyridyl) methylamine in a PVC-Membrane Sensor and Characterization of the Membrane Properties. *Chem Cent J.* 2012;6(1):1-6.
55. Duarte AF, Ferreira S, Oliveira R, Domingues FC. Effect of coriander oil (*Coriandrum sativum*) on planktonic and biofilm cells of *Acinetobacter baumannii*. *Nat Prod Commun.* 2013;8(5):1934578X1300800532.
56. Almessiere MA, Slimani Y, Rehman S, Khan FA, Sertkol M, Baykal A. Green synthesis of Nd substituted Co-Ni nanospinel ferrites: a structural, magnetic, and antibacterial/anticancer investigation. *J Phys Appl Phys.* 2021;55:055002.
57. Al-Jameel SS, Rehman S, Almessiere MA, et al. Anti-microbial and anti-cancer activities of Mn_{0.5}Zn_{0.5}DyxFe_{2-x}O₄ (x ≤ 0.1) nanoparticles. *Artif Cell Nanomed Biotechnol*;49(2021):493-499.
58. Almessiere M, Slimani Y, Auwal I, et al. Biosynthesis effect of Moringa oleifera leaf extract on structural and magnetic properties of Zn doped Ca-Mg nano-spinel ferrites. *Arab J Chem.* 2021;14:103261.