

Original Article

# Evaluation of Phytochemicals and Antibacterial Usefulness of Citrus Limon & Cicer Arietinum in Synergistic Effect with Antibiotics Against Clinically Important Bacteria

Aamir Sattar<sup>1\*</sup>, Afshan Syed Abbas<sup>2</sup>, Sajida Naseem<sup>3</sup>, Madiha Mumtaz<sup>4</sup>, Asad Shabbir<sup>5</sup>, Saima Shahzad Mirza<sup>2</sup>

<sup>1</sup>University of Education Lower Mall Campus Lahore, Pakistan.

<sup>2</sup>University of Education Bank Road Campus Lahore, Pakistan.

<sup>3</sup>University of Education, Township Campus Lahore, Pakistan.

<sup>4</sup>Khawaja Muhammad Safdar Medical College Sialkot, Pakistan.

<sup>5</sup>University of Sialkot, Pakistan.

\*Corresponding Author: Aamir Sattar; Email: aaamirsattar@gmail.com

Conflict of Interest: None.

Sattar A., et al. (2024). 4(1): DOI: <https://doi.org/10.61919/jhrr.v4i1.174>

## ABSTRACT

**Background:** The emergence of antibiotic resistance necessitates exploring alternative treatments, including natural plant extracts. Lemon juice (Citrus limon) and chickpea (Cicer arietinum) seed aqueous extract, known for their medicinal properties, are investigated for their potential antibacterial effects.

**Objective:** This study aimed to assess the antibacterial efficacy of lemon juice and chickpea seed aqueous extract, individually and in combination with antibiotics, against various pathogenic bacterial strains.

**Methods:** Aqueous and ethanolic extracts of C. limon and C. arietinum were prepared. Antibacterial activity against Escherichia coli, Klebsiella pneumoniae, Staphylococcus aureus, and Pseudomonas aeruginosa, sourced from the University of Education, Lahore, was assessed using the disc diffusion method. Extract concentrations of 5%, 10%, 20%, and 40% were tested. Phytochemical analysis identified active compounds in the extracts. Antibacterial efficacy was quantified by measuring inhibition zones in millimeters.

**Results:** Lemon juice showed dose-dependent antibacterial activity with inhibition zones up to  $19.8 \pm 0.088$  mm at 40% concentration. Chickpea extract was effective against S. aureus with a maximum inhibition of  $14 \pm 0.577$  mm at 40%. Combined treatments enhanced antibacterial effects, particularly against S. aureus, with a combination of lemon juice, chickpea extract, and antibiotics yielding an inhibition zone of  $20.1 \pm 0.058$  mm at the highest concentration. In contrast, chickpea extract alone showed no activity against E. coli, K. pneumoniae, or P. aeruginosa.

**Conclusion:** The study confirms the significant antibacterial potential of lemon juice across multiple bacterial strains and the specific efficacy of chickpea extract against S. aureus. The combination of these extracts with antibiotics suggests a promising avenue for augmenting antibacterial treatment strategies.

**Keywords:** Antibacterial Activity, Citrus limon, Cicer arietinum, Plant Extracts, Antibiotic Resistance, Alternative Therapies.

## INTRODUCTION

Antibiotics, a cornerstone in the battle against bacterial infections, were first discovered by Alexander Fleming in 1928, marking a revolution in medical science and saving countless lives (1). Gerhard Domagk's discovery of the antiseptic Prontosil in 1939 further advanced this field, earning him the Nobel Prize (2, 3). However, the initial triumph was soon tempered by the emergence of antibiotic resistance, a phenomenon first observed in E. coli and S. aureus, which showed resistance to penicillin shortly after its discovery (4, 5). Today, antibiotic resistance poses a significant challenge in modern medicine, with an increasing number of pathogens demonstrating reduced sensitivity to commonly used antibiotics (6). This alarming trend of harmful resistant microbes has been escalating over the years (7).

Parallel to the developments in synthetic medicine, the traditional use of plant extracts for disease treatment has been well-documented, with numerous plants identified for their medicinal properties (9). Among these, the Citrus genus, a member of the Rutaceae family, holds a prominent place in traditional medicine. Citrus limon, in particular, is celebrated not only for its delightful

taste but also for its array of therapeutic properties. It offers a rich source of essential nutrients and bioactive compounds, including vitamins and secondary metabolites, which contribute to health enhancement and disease prevention (15). Citrus Limon is noted for its diverse pharmacological activities, including its antibacterial and anticancer properties, attributed to its content of alkaloids, vitamins C, flavonoids, phenols, steroids, reducing sugar, and other antioxidants (16, 17, 18, 26). Genomic studies reveal that *C. limon* originated as a hybrid between the acrid orange and lime, with its exact origins believed to be in India.

*Cicer arietinum*, commonly known as chickpea, is a staple in diets across various regions of the world and is recognized for its high content of proteins, fibers, minerals, and vitamins (10). Notable for having the highest oil content among dry pulses (3-10%), *C. arietinum* is also free from anti-diet elements and rich in carotenoids, such as beta-carotene (11, 12). Research by Dr. Maryam M. Elmi & Ozcelik et al. highlighted the antibacterial and antifungal properties of *C. arietinum*, suggesting its potential use in controlling microbial growth (13, 14).

The present study focuses on evaluating the antibacterial properties of aqueous and ethanolic extracts of *C. limon* and *C. arietinum* against commonly resistant pathogenic bacterial strains like *E. coli*, *P. aeruginosa*, *S. aureus*, and *K. pneumoniae*. Additionally, the study delves into analyzing the phytochemical constituents of these extracts, identifying the specific components responsible for their antimicrobial activity. This research contributes to the growing body of knowledge on alternative antimicrobial strategies, particularly in the context of increasing antibiotic resistance, and underscores the potential of these plant extracts in developing new therapeutic agents.

## MATERIAL AND METHODS

In this study, we examined the antibacterial properties of Citrus limon (*C. limon*) and Cicer arietinum (*C. arietinum*) against several resistant bacterial strains. The methodology encompassed the preparation of plant extracts, their phytochemical analysis, and the assessment of their antibacterial activity.

Bacterial cultures of *Escherichia coli*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, and *Pseudomonas aeruginosa* were obtained from the Zoological Lab at the University of Education, Lower Mall Campus, Lahore, Pakistan. Fresh samples of *C. limon* were procured from a local market, and *C. arietinum* seeds were acquired from the same source.

The aqueous extract of *C. arietinum* was prepared by first cleaning the seeds to remove dirt and other materials, followed by air-drying at room temperature. The dried seeds were then ground into a fine powder. Following the method of Basri and Fan (21), a 10% solution was prepared by mixing 50 grams of the powdered seeds with 500 mL of distilled water. This mixture was agitated continuously for 24 hours using a magnetic stirrer and then filtered through Whatman filter paper no. 1. The extract was pasteurized using a 0.45µm filter paper to ensure sterility and the absence of germs.

For the preparation of *C. limon* pulp, fresh fruits were halved and squeezed to obtain the pulp. The antibacterial activity of both *C. arietinum* and *C. limon* extracts was assessed using the disc diffusion method. Filter paper discs of approximately 5 mm in diameter were placed on petri plates. Then, 10 µL of the fresh aqueous and ethanolic extracts of *C. arietinum* and *C. limon* pulp were applied to each of the bacterial species. Amoxicillin was used as a positive control, while the respective solvents (distilled water and ethanol) served as negative controls (21). The experimental conditions, including time gap, dosage concentration, temperature range, and environmental exposure, were meticulously controlled. The inhibition zones were measured after 24 hours using Vernier calipers.

Phytochemical analysis was conducted to identify the active constituents in the extracts. Various tests were performed to detect the presence of tannins, glycosides, proteins, cardiac glycosides, steroids, phytosterols, phenols, flavonoids, carbohydrates, alkaloids, and saponins using standard methods (1, 9, 10, 11, 15, 17, 18, 21). The specific procedures included heating, addition of reagents like ferric chloride, Fehling's solution, Millon's reagent, Ninhydrin, sulfuric acid, and observing color changes or precipitate formation. All methods were carried out following the relevant guidelines and regulations. Ethical approval was obtained from the institutional review board. Data collection was systematic, ensuring accuracy and reliability. The data analysis involved quantitative measurements of inhibition zones and qualitative assessments of phytochemical tests. The results were then statistically analyzed to determine the efficacy of the plant extracts against the bacterial strains. This comprehensive approach provided a thorough understanding of the antibacterial potential and chemical composition of *C. limon* and *C. arietinum*.

## RESULTS

The results of this study, as presented in Tables 1 to 4, reveal insightful observations regarding the antibacterial effects of various treatments against different bacterial strains.

Table 1 MEAN (mm)  $\pm$ SEM values of antibacterial effect of various concentrations of (Anti) Positive control, (L.J) Lemon juice, (C.E) Chickpea aqueous extract (D.W) Negative control, on *P. aeruginosa*

| Treatments      | 5%              | 10%             | 20%              | 40%              |
|-----------------|-----------------|-----------------|------------------|------------------|
| Anti            | 1.1 $\pm$ 0.058 | 4.1 $\pm$ 0.058 | 9.03 $\pm$ 0.033 | 19.8 $\pm$ 0.088 |
| L.J             | 0.9 $\pm$ 0.058 | 3.1 $\pm$ 0.058 | 7.9 $\pm$ 0.058  | 18.8 $\pm$ 0.088 |
| C.E             | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  |
| D.W             | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  |
| Anti + C.E      | 2.1 $\pm$ 0.088 | 5.1 $\pm$ 0.058 | 8.1 $\pm$ 0.058  | 17.1 $\pm$ 0.058 |
| Anti + L.J      | 2.1 $\pm$ 0.058 | 7.1 $\pm$ 0.058 | 16.1 $\pm$ 0.058 | 21.1 $\pm$ 0.058 |
| L.J + C.E       | 1.9 $\pm$ 0.058 | 4.9 $\pm$ 0.058 | 8.96 $\pm$ 0.088 | 14.1 $\pm$ 0.033 |
| Anti + L.J+ C.E | 2.1 $\pm$ 0.058 | 6.1 $\pm$ 0.058 | 11.1 $\pm$ 0.058 | 18.1 $\pm$ 0.058 |

In Table 1, the antibacterial effect of treatments against *P. aeruginosa* is detailed. The positive control (Anti) demonstrated a dose-dependent increase in efficacy, with mean inhibition zones ranging from 1.1 $\pm$ 0.058 mm at 5% concentration to 19.8 $\pm$ 0.088 mm at 40%. Lemon juice (L.J) also showed a similar trend but with slightly lower effectiveness, reaching an inhibition zone of 18.8 $\pm$ 0.088 mm at the highest concentration. Notably, chickpea aqueous extract (C.E) and the negative control (D.W) exhibited no antibacterial activity across all concentrations. The combinations of treatments, particularly Anti + L.J and L.J + C.E, indicated enhanced antibacterial effects compared to individual treatments, with Anti + L.J reaching an inhibition zone of 21.1 $\pm$ 0.058 mm at 40%.

Table 2 MEAN (mm)  $\pm$ SEM values of antibacterial effect of various concentrations of (Anti) Positive control, (L.J) Lemon juice, (C.E) Chickpea aqueous extract and (D.W) Negative control, on *K. pneumoniae*.

| Treatments     | 5%              | 10%             | 20%              | 40%              |
|----------------|-----------------|-----------------|------------------|------------------|
| Anti           | 2.0 $\pm$ 0.058 | 7.1 $\pm$ 0.058 | 11.0 $\pm$ 0.067 | 19.1 $\pm$ 0.058 |
| L.J            | 0.9 $\pm$ 0.033 | 6.1 $\pm$ 0.088 | 10.1 $\pm$ 0.058 | 18.0 $\pm$ 0.067 |
| C.E            | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  |
| D.W            | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  |
| Anti + C.E     | 1.1 $\pm$ 0.033 | 5.9 $\pm$ 0.058 | 9.1 $\pm$ 0.058  | 15.1 $\pm$ 0.058 |
| Anti + L.J     | 3.0 $\pm$ 0.058 | 8.1 $\pm$ 0.058 | 15.0 $\pm$ 0.033 | 21.0 $\pm$ 0.033 |
| L.J + C.E      | 1.1 $\pm$ 0.058 | 6.0 $\pm$ 0.033 | 8.9 $\pm$ 0.058  | 12.1 $\pm$ 0.058 |
| Anti+ L.J+ C.E | 2.1 $\pm$ 0.058 | 6.1 $\pm$ 0.058 | 11.1 $\pm$ 0.058 | 18.1 $\pm$ 0.058 |

In Table 2, the focus shifts to *K. pneumoniae*. Here again, the positive control showed a consistent increase in antibacterial activity with increasing concentrations, peaking at 19.1 $\pm$ 0.058 mm at 40%. Lemon juice's efficacy was notable, especially at higher concentrations, reaching 18.0 $\pm$ 0.067 mm at 40%. The combination treatments, particularly Anti + L.J, demonstrated a synergistic effect, achieving an inhibition zone of 21.0 $\pm$ 0.033 mm at 40%.

Table 3 MEAN (mm)  $\pm$ SEM values of antibacterial effect of various concentrations of (Anti) Positive control, (L.J) Lemon juice, (C.E) Chickpea aqueous extract and (D.W) Negative control, on *E.coli*.

| Treatments      | 5%              | 10%             | 20%              | 40%              |
|-----------------|-----------------|-----------------|------------------|------------------|
| Anti            | 1.0 $\pm$ 0.066 | 5.3 $\pm$ 0.033 | 7.6 $\pm$ 0.33   | 18.3 $\pm$ 0.333 |
| L.J             | 1.1 $\pm$ 0.057 | 3.9 $\pm$ 0.067 | 7.3 $\pm$ 0.33   | 17.5 $\pm$ 0.280 |
| C.E             | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  |
| D.W             | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000 | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  |
| Anti + C.E      | 1.0 $\pm$ 0.033 | 4.1 $\pm$ 0.057 | 7.76 $\pm$ 0.14  | 14.5 $\pm$ 0.288 |
| Anti + L.J      | 3 $\pm$ 0.057   | 6.4 $\pm$ 0.200 | 14.0 $\pm$ 0.033 | 19.8 $\pm$ 0.155 |
| L.J + C.E       | 0.9 $\pm$ 0.033 | 5.0 $\pm$ 0.057 | 9.0 $\pm$ 0.033  | 14.1 $\pm$ 0.057 |
| Anti + L.J+ C.E | 2.1 $\pm$ 0.088 | 6.7 $\pm$ 0.140 | 8.7 $\pm$ 0.33   | 17.1 $\pm$ 0.08  |

Table 4 MEAN (mm)  $\pm$ SEM values of antibacterial effect of various concentrations of (Anti) Positive control, (L.J) Lemon juice, (C.E) Chickpea aqueous extract (D.W) Negative control on *S. aureus*.

| Treatments     | 5%                         | 10%              | 20%              | 40%               |
|----------------|----------------------------|------------------|------------------|-------------------|
| Anti           | 3.0 $\pm$ 0.0338.133 $\pm$ | 8.1 $\pm$ 0.088  | 12.5 $\pm$ 0.289 | 17.4 $\pm$ 0.200  |
| L.J            | 2.1 $\pm$ 0.888            | 7.0 $\pm$ 0.033  | 11.6 $\pm$ 0.33  | 15.73 $\pm$ 0.144 |
| C.E            | 0.0 $\pm$ 0.000            | 2.8 $\pm$ 0.088  | 7.8 $\pm$ 0.133  | 14 $\pm$ 0.577    |
| D.W            | 0 $\pm$ 0.00               | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000  | 0.0 $\pm$ 0.000   |
| Anti + C.E     | 3.9 $\pm$ 0.067            | 8.93 $\pm$ 0.067 | 14.3 $\pm$ 0.153 | 18.1 $\pm$ 0.058  |
| Anti + L.J     | 1.0 $\pm$ 0.000            | 6.86 $\pm$ 0.133 | 11.9 $\pm$ 0.033 | 19.03 $\pm$ 0.033 |
| L.J + C.E      | 0.0 $\pm$ 0.000            | 7.1 $\pm$ 0.588  | 13.1 $\pm$ 0.053 | 19.1 $\pm$ 0.058  |
| Anti+ L.J+ C.E | 2.0 $\pm$ 0.033            | 7.2 $\pm$ 0.100  | 12.1 $\pm$ 0.53  | 20.1 $\pm$ 0.058  |

The results against *E. coli*, presented in Table 3, mirrored the trends observed in previous tables. The positive control showed progressive effectiveness, with a maximum zone of inhibition of 18.3 $\pm$ 0.333 mm at 40%. The combination of Anti + L.J was particularly effective, exhibiting an inhibition zone of 19.8 $\pm$ 0.155 mm at the highest concentration. Interestingly, the combination of L.J + C.E also showed considerable activity, with an inhibition zone of 14.1 $\pm$ 0.057 mm at 40%.

Table 4 presents the results against *S. aureus*. The positive control maintained its pattern of increased efficacy with higher concentrations, reaching an inhibition zone of 17.4 $\pm$ 0.200 mm at 40%. The chickpea aqueous extract displayed activity at higher concentrations, with an inhibition zone of 14 $\pm$ 0.577 mm at 40%, a notable finding given its lack of activity against other bacterial strains. The combinations, particularly L.J + C.E and Anti + L.J + C.E, exhibited enhanced antibacterial effects, with the latter achieving the largest zone of inhibition of 20.1 $\pm$ 0.058 mm at 40%.

Overall, the results underscore the potential of lemon juice and chickpea aqueous extract, both individually and in combination with the positive control, in exerting antibacterial effects against various bacterial strains. The synergistic effects observed in the combination treatments are particularly promising, suggesting a potential avenue for enhancing antibacterial efficacy, especially in the context of increasing antibiotic resistance.

## DISCUSSION

The study explored the antibacterial properties of lemon juice and chickpea seed aqueous extract, both individually and in combination with antibiotics, against various bacterial strains. The findings revealed that lemon juice, containing vital substances such as antioxidants, flavonoids, phenols, steroids, reducing sugars, and alkaloids, exhibited significant antimicrobial effects. This aligns with previous research indicating lemon's curative potential (16, 22), and its effectiveness against diarrhea-causing bacteria due to its rich content of potassium, vitamin C, alkaloids, and citric acid (16, 23, 24, 25). Mathai K. and colleagues also corroborated lemon juice's antibacterial potential, particularly against *S. mutants* (26).

The antibacterial mechanism of flavonoids, a key component in lemon, involves inhibiting nucleic acid synthesis, disrupting cytoplasmic membrane functions, affecting energy metabolism, and impacting biofilm synthesis and membrane permeability (32, 33). Flavonols and phenolic acids, in particular, demonstrate substantial antibacterial action, inhibiting bacterial virulence factors such as toxins and enzymes, and enhancing the efficacy of antibiotics (34).

Conversely, chickpea seed aqueous extract displayed limited antibacterial activity. Previous studies by Mallikarjuna et al. (2007) found chickpea extract effective against fungi and identified antifungal peptides cicerin and arietin in chickpea seeds (27, 31). While legumes like chickpeas are known for their antibacterial effects against *Bacillus cereus* and antioxidant potential (28), the current study found chickpea extracts effective against *S. aureus* but not against *K. pneumoniae*, *P. aeruginosa*, or *E. coli*.

The study's distinctiveness lies in its exploration of the combined effects of antibiotics, lemon juice, and chickpea seed aqueous extracts. Notably, the synergistic combination of lemon juice and antibiotics showed enhanced antibacterial activity compared to antibiotics alone. In contrast, the combination of chickpea aqueous extract and antibiotics was more effective only against *S. aureus*. However, the study had limitations. The scope was confined to in vitro conditions, which may not fully replicate in vivo environments. Future research should consider in vivo studies to validate these findings. Additionally, the specific mechanisms through which these extracts exert their antibacterial effects, particularly in combination treatments, warrant further investigation.

## CONCLUSION

In conclusion, lemon juice demonstrated broad-spectrum antibacterial properties, while chickpea seed aqueous extract showed limited activity. The combinations, especially of lemon juice and antibiotics, presented a promising avenue for enhancing

antibacterial efficacy. This study underscores the potential of using plant sources in developing alternative or complementary antibacterial therapies.

Funding for this study was generously provided by the Zoology Department, University of Education Lahore, Pakistan.

Acknowledgements are due to Dr. Zahid Iqbal, Dr. Anwar Malik, Dr. Saleema Bashir, Miss Razia Bashir, Miss Nusrat Majeed, and Miss Saman Javeed for their invaluable contributions and support.

## REFERENCES

1. Perry, C. M. (2021). Antibiotic Resistance Crisis Spurring Phage Therapy Research.
2. Pegadraj, H., Thomas, J. A., Mutha, J. A., & Reshmi, R. RECAPITULATION OF ANTIBIOTIC RESISTANCE AND MRSA.
3. Tan, S.Y. and Tatsumura, Y., 2015. Alexander Fleming (1881–1955): discoverer of penicillin. Singapore medical journal, 56(7), p.366.
4. Abraham EP, Chain E. 1988. An enzyme from bacteria able to destroy penicillin. 1940. Rev Infect Dis 10:677-8
5. Rammelkamp CH, Maxon T. 1942. Resistance of Staphylococcus aureus to the Action of Penicillin. Proceedings of the Society for Experimental Biology and Medicine 51:386-389.
6. Chen, Z., Li, Y., Peng, Y., Ye, C., & Zhang, S. (2021). Effects of antibiotics on hydrolase activity and structure of microbial community during aerobic co-composting of food waste with sewage sludge. Bioresource Technology, 321, 124506
7. Fair, R.J. and Tor, Y., 2014. Antibiotics and bacterial resistance in the 21st century. Perspectives in medicinal chemistry, 6, pp.PMC-S14459.
8. Miranda, C., Silva, V., Igrejas, G., & Poeta, P. (2021). Impact of European pet antibiotic use on enterococci and staphylococci antimicrobial resistance and human health. Future Microbiology, 16(3), 185-203.
9. Okeke, I. N., Laxminarayan, Bhutta, Z. A., Duse, Jenkins, P., O' Brien, T. F. & Klugman, K. P. 2005
10. Moutassem, D., Belabid, L. and Bellik, Y., 2020. Efficiency of secondary metabolites produced by Trichoderma spp. in the biological control of Fusarium wilt in chickpea. Journal of Crop Protection, 9(2), pp.217-231.
11. De Almeida Costa, G. E., da Silva Queiroz-Monici, K., Reis, S. M. P. M. & de Oliveira, A. C. 2006
12. Mallikarjuna, N. N. & Varma, R. S. 2007; Abbo, S., Lev Yadun, S. & Gopher, A. 2011
13. Barari, L., Mosavi, N., & Asgharpour, F. (2015). Antibacterial and antifungal effect of chickpea (cicer arietinum) aqueous seed extract. World Journal of Pharmacy and Pharmaceutical Sciences, 4(10), 336-346.
14. Kan, Z., Jaiswal, B.S., Stinson, J., Janakiraman, V., Bhatt, D., Stern, H.M., Yue, P., Haverty, P.M., Bourgon, R., Zheng, J. and Moorhead, M., 2010. Diverse somatic mutation patterns and pathway alterations in human cancers. Nature, 466(7308), pp.869-873.
15. Jouda, M.M., 2013. The antibacterial effect of some medicinal plant extracts and their synergistic effect with antibiotic and non-antibiotic drugs. The Antibacterial Effect of Some Medicinal Plant Extracts and their Synergistic Effect with Antibiotic and Non-antibiotic Drugs.
16. Mohanapriya, M., Ramaswamy, L. and Rajendran, R., 2013. Health and medicinal properties of lemon (Citrus limonum). International Journal Of Ayurvedic And Herbal Medicine, 3(1), pp.1095-1100.
17. Ekawati, E.R. and Darmanto, W., 2019. Lemon (citrus limon) juice has antibacterial potential against diarrhea-causing pathogen. In IOP Conference Series: Earth and Environmental Science (Vol. 217, No. 1, p. 012023). IOP Publishing.
18. Makni, M., Jemai, R., Kriaa, W., Chtourou, Y. and Fetoui, H., 2018. Citrus limon from Tunisia: Phytochemical and physicochemical properties and biological activities. BioMed research international, 2018.
19. Gibbons, S. (2004). Anti-staphylococcal activity of natural plants. Natural Products Reports, 21, 263-277.
20. Coutinho, H.D., Costa, J.G., Lima, E.O., Falcão-Silva, V.S. and Júnior, J.P.S., 2009. Herbal therapy associated with antibiotic therapy: potentiation of the antibiotic activity against methicillin-resistant Staphylococcus aureus by Turnera ulmifolia L. BMC complementary and alternative medicine, 9(1), p.13.
21. D.F.BasriandS.H.Fan, "The potential of aqueous and acetone extracts of galls of Quercus infectoria as antibacterial agents," Indian Journal of Pharmacology, vol.37, no.1, pp.26–29, 2005.
22. Viuda-Martos, M., Ruiz-Navajas, Y., Fernández-López, J., & Pérez-Álvarez, J. (2008). Antifungal activity of lemon (Citrus lemon L.), mandarin (Citrus reticulata L.), grapefruit (Citrus paradisi L.) and orange (Citrus sinensis L.) essential oils. Food control, 19(12), 1130-1138.
23. Yadav, G., & Meena, M. (2021). Bioprospecting of endophytes in medicinal plants of Thar Desert: An attractive resource for biopharmaceuticals. Biotechnology Reports, e00629.
24. Gibbons, S. (2004). Anti-staphylococcal activity of natural plants. Natural Products Reports, 21, 263-277.
25. Coutinho, H. D., Costa, J. G., Falcão-Silva, V. S., Lima, E. O. & Siqueira Júnior, J. P. 2008

26. Thilakchand, K. R., Mathai, R. T., Simon, P., Ravi, R. T., Baliga-Rao, M. P., & Baliga, M. S. (2013). Hepatoprotective properties of the Indian gooseberry (*Emblica officinalis* Gaertn): a review. *Food & function*, 4(10), 1431-1441.
27. Guo, D., Liu, W., Zhang, X., Zhao, M., Zhu, B., Hou, T. and He, H., 2019. Duck Egg White-Derived Peptide VSEE (Val-Ser-Glu-Glu) Regulates Bone and Lipid Metabolisms by Wnt/ $\beta$ -Catenin Signaling Pathway and Intestinal Microbiota. *Molecular Nutrition & Food Research*, 63(24), p.1900525.
28. Mejri, F., Selmi, S., Martins, A., Baati, T., Chaabane, H., Njim, L., Serralheiro, M.L., Rauter, A.P. and Hosni, K., 2018. Broad bean (*Vicia faba* L.) pods: A rich source of bioactive ingredients with antimicrobial, antioxidant, enzyme inhibitory, anti-diabetic and health-promoting properties. *Food & function*, 9(4), pp.2051-2069.
29. Saquib, S.A., AlQahtani, N.A., Ahmad, I., Kader, M.A., Al Shahrani, S.S. and Asiri, E.A., 2019. Evaluation and comparison of antibacterial efficacy of herbal extracts in combination with antibiotics on periodontal pathobionts: an in vitro microbiological study. *Antibiotics*, 8(3), p.89.
30. Olajuyigbe, O.O. and Afolayan, A.J., 2013. Evaluation of combination effects of ethanolic extract of *Ziziphus mucronata* Willd. subsp. *mucronata* Willd. and antibiotics against clinically important bacteria. *The Scientific World Journal*, 2013.
31. Mathai, K., Anand, S., Aravind, A., Dinatius, P., Krishnan, A.V. and Mathai, M., 2017. Antimicrobial Effect of Ginger, Garlic, Honey, and Lemon Extracts on *Streptococcus mutans*. *The journal of contemporary dental practice*, 18(11), pp.1004-1008.
32. (Xie, Y., Yang, W., Tang, F., Chen, X., & Ren, L. (2015). Antibacterial activities of flavonoids: structure-activity relationship and mechanism. *Current medicinal chemistry*, 22(1), 132-149.)
33. Naqvi, S. A. R., Nadeem, S., Komal, S., Naqvi, S. A. A., Mubarik, M. S., Qureshi, S. Y., ... & Aslam, N. (2019). Antioxidants: Natural Antibiotics. In *Antioxidants*. IntechOpen.).
34. (Mikłasińska-Majdanik, M., Kępa, M., Wojtyczka, R. D., Idzik, D., & Wąsik, T. J. (2018). Phenolic compounds diminish antibiotic resistance of *Staphylococcus aureus* clinical strains. *International journal of environmental research and public health*, 15(10), 2321.)