

Original Article

Elemental composition of Talang Queenfish (*Scomberoides commersonnianus* Lacépède, 1801) in relation to body size and condition factor from Pakistan

Composição elementar de Talang Queenfish (*Scomberoides commersonnianus* Lacépède, 1801) em relação ao tamanho do corpo e fator de condição do Paquistão

S. M. Azam^{a,b} , M. Naeem^{a*} , N. Ahmad^c , R. Yaqoob^b , M. Khalid^a  and V. Lal^a 

^aBahauddin Zakariya University, Institute of Zoology, Multan, Pakistan

^bUniversity of Education, Department of Zoology, Division of Science & Technology, Lahore, Pakistan

^cUniversity of Jhang, Department of Zoology, Punjab, Pakistan

Abstract

Fish is thought to be the most appropriate for indication of toxicity in water. So, purpose of present study was to determine the concentrations of few metals (Na, K, Ca, Mg, Fe, Cd, Pb and Zn) in relation to the body size and condition factor in *Scomberoides commersonnianus*. A total of 73 samples of marine fish, *S. commersonnianus* with various sizes that ranged from 93.4–1180 g of body weight and total length 20.5–56.9 cm were randomly collected from the Arabian Sea of Karachi Pakistan, for examination of elemental composition. The concentration of Calcium found maximum while Cadmium found minimum in *S. commersonnianus*. Concentrations of Zn showed highly significant ($P < 0.001$) negative, while Na and Ca exhibited strong positive correlation with the fish size. Correlations of Pb, K and Mg were found least significant ($P < 0.05$) with the wet weight and total length of *S. commersonnianus*. Whereas, Fe and Cd were remained insignificantly correlated ($P > 0.05$) with the fish size. Condition factor showed highly significant correlation ($P < 0.001$) with all studied metals except cadmium and lead which were found insignificantly correlated ($P > 0.05$) with condition factor of *S. commersonnianus*.

Keywords: marine fish, *Scomberoides commersonnianus*, metal concentrations, fish size, condition factor.

Resumo

O peixe é considerado o mais adequado para indicação de toxicidade na água. Assim, o objetivo do presente estudo foi determinar as concentrações de alguns metais (Na, K, Ca, Mg, Fe, Cd, Pb e Zn) em relação ao tamanho corporal e fator de condição em *Scomberoides commersonnianus*. Foi coletado um total de 73 amostras de peixes marinhos, *S. commersonnianus*, com tamanhos que variaram de 93,4 a 1.180 g de peso corporal e comprimento total de 20,5 a 56,9 cm, aleatoriamente do Mar Árábico de Karachi, Paquistão, para exame de composição elementar. A concentração de Ca foi máxima, enquanto a de Cd foi mínima em *S. commersonnianus*. As concentrações de Zn mostraram-se altamente significativas negativas ($P < 0,001$), enquanto Na e Ca apresentaram forte correlação positiva com o tamanho dos peixes. As correlações de Pb, K e Mg foram menos significativas ($P < 0,05$) com o peso úmido e o comprimento total de *S. commersonnianus*. Já o Fe e o Cd permaneceram pouco correlacionados ($P > 0,05$) com o tamanho dos peixes. O fator de condição apresentou correlação altamente significativa ($P < 0,001$) com todos os metais estudados, exceto Cd e Pb, que foram encontrados correlacionados insignificamente ($P > 0,05$) com o fator de condição de *S. commersonnianus*.

Palavras-chave: peixes marinhos, *Scomberoides commersonnianus*, concentrações de metais, tamanho do peixe, fator de condição.

1. Introduction

Fishes are distinct from other species in that they have the capacity to absorb minerals from both their diet and the water through their skin and gills (Abbas et al., 2021). Fish is thought to be the most accurate indicator of water toxicity (Barse et al., 2007; Nasir and Al-Najar, 2015). When fishes are exposed to higher levels of toxic metals, these

harmful contaminants are taken up by the fish body largely, affecting the human body's health directly because of eating polluted feed (Nasir and Al-Najar, 2015). As a result, the contamination of fish with these trace elements are being given considerable attention (Coetzee, 1998). The pollutant may enter in the body of fish and absorbed in the kidney and

*e-mail: dr_naeembzu@yahoo.com

Received: March 29, 2022 – Accepted: May 21, 2022



This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gills, eventually become the parts of animal's body mass (Gasperi et al., 2018). Toxicity in the body of the fish is affected by its age, size, and sexual behavior. Metal evaluations in fish tissue are necessary to ensure that metal deposition in fish does not pass on to fish consumers, such as humans (Rahman et al., 2012). Once pollutants are released into the water, it does not stay in that location but spreads to other parts of water (Everaert et al., 2018). Heavy metals have an initial influence on the fish at a cellular or tissue level, but later on, major changes in fish physiology and to some extent, its behavior have been recorded (Asegbeloyi et al., 2010). Several elements such as Copper, Lead, Cadmium, Zinc, and Iron were found in the food and water (Staniskiene et al., 2009). The concentration responses of these elements in relation to their presence and harmful effects on human and other fish have been studied (Koelmans et al., 2017). Some organisms have great capacity for absorbing heavy elements in their bodies, whereas others have none or a low potential for doing so. Some fish absorb much from food, and the harmful substances become a part of their bodies as a result of their excessive consumption (Eisler, 1987). Concentration of hazardous heavy metal components in freshwater fish in Pakistan suggest that rapid mechanical and agricultural improvement in the past has posed significant threats to the country's water resources (Ahmad et al., 2021).

This research was performed to examine the relationship of fish size and condition factor with some mineral elements in marine Talang Queenfish (*Scomberoides commersonnianus*).

2. Materials and Methods

A total of 73 samples of a marine fish, *S. commersonnianus* with various sizes that ranged from 93.4–1180g of body weight and total length 20.5–56.9 cm were randomly collected from the Arabian Sea Karachi, Sindh, Pakistan, for examination of elemental composition. In plastic bags, these samples were delivered to Fish Research Lab, Institute of Zoology, Bahauddin Zakariya University, Multan, Pakistan. These fish samples were homogenized into fine powder after drying in the oven. One gram of powdered-sample was taken from each sample for ashing and kept on 500°C in muffle furnace for 1 day. Ashes were then dissolved into the 70% HNO₃ (10 ml); heated at 80 to 100°C on hot plate and diluted upto 25ml with deionized water. These samples were run on Atomic Absorption Spectrophotometer (Agilent, AA 200) at Institute of Zoology, BZU, Multan, Pakistan, for quantification of Zn, Cd, Fe, Ca, Na, K, Pb and Mg concentration in samples. Repeatedly calibrations were set on instrument after every ten metal samples during operation.

To access and evaluate the relationship of these metal concentrations with the body size and condition factor of fish samples, regression analysis had been performed with the help of following Equation 1:

$$\text{Log } Y = \text{Log } a + b \text{ Log } X \quad (1)$$

Here:

Y stand for metal concentration ($\mu\text{g g}^{-1}$); X = Fish length (cm)/Weight (g)/Condition Factor; *a* for constant and *b* for slope.

At $P < 0.001$, $P < 0.01$ and $P < 0.05$, correlation coefficients had been considered as significant. Using *b* value, log data was used to analyze and explain the fish's allometric metal concentration pattern.

Multiple regression studies to access the relationships between fish size and metal concentration were calculated through following Equation 2:

$$Y = a + b_1 WW + b_2 TL \quad (2)$$

Where Y for metal concentration, and *b*₁, *b*₂ representing slope values, *a* is constant, WW for wet body weight and TL for total length. For statistical analyses, Microsoft Excel 2016 and MiniTab 18 were used.

3. Results

3.1. Determination of metal concentrations in wet and dry body weight of *S. commersonnianus*.

The concentration of Calcium found to be maximum while Cadmium found to be minimum in *S. commersonnianus*. Mean ($\pm$ SD) concentrations of Zn, Fe, Cd, Pb, Na, K, Ca and Mg had been found 41.60 ± 17.20 , 59.32 ± 53.06 , 0.04 ± 0.09 , 0.20 ± 0.29 , 589.16 ± 247.20 , 896.62 ± 268.85 , 1577.90 ± 569.04 and $307.12 \pm 118.73 \mu\text{g/g}$, respectively, in wet body weight of *S. commersonnianus*. Descriptive statistics of metal concentration in wet and dry body weight of *S. commersonnianus* are shown in Table 1.

3.2. Correlation of various metals with total length and wet weight of *S. commersonnianus*

Different metals were examined in correlation with wet weight and total length of *S. commersonnianus*, and regression parameters are represented in Table 2 and 3, respectively. Zn showed highly significant negative correlation ($P < 0.001$), while Na and Ca represented strong positive correlation ($P < 0.001$) with fish size (wet weight and total length). Correlations of Pb, K, Mg were found least significant ($P < 0.05$) with the total length and wet weight of *S. commersonnianus*. However, concentrations of Fe and Cd were remained insignificantly correlated ($P > 0.05$) with the fish size.

Regression parameters of condition factor with metal concentration in wet body weight for *S. commersonnianus* are presented in Table 4. K, Zn, Na, Mg, Ca and Fe exhibited highly significant positive correlation with total body length ($P < 0.001$) with values ranging from 0.441 to 0.711, while Cd and Pb exhibited non-significant positive correlation ($P > 0.05$) in *S. commersonnianus*.

Multiple regression parameters of total length and wet weight with concentration of metal for *S. commersonnianus* are presented in Table 5. Concentration of K, Na, Mg, Ca, Cd and Fe exhibited highly significant positive correlation ($P < 0.001$) with size of *S. commersonnianus* with multiple regression values ranging from 0.456 to 0.886; Zn showed least significant positive correlation ($P > 0.05$) with value ($r = 0.285$); while, Pb showed insignificant positive correlation ($P < 0.05$). Results of multiple regression analysis

Table 1. Descriptive analysis of element concentration (µg/g) in wet and dry body weight of wild *Scomberoides commersonnianus* (n = 73).

Elements	Range		Mean±SD	
	wet weight	dry weight	wet weight	dry weight
Fe	9.58-86.14	34.34-422.50	41.60±17.20	162.74±75.14
Zn	0.06-199.10	0.24-734.19	59.32±53.06	237.57±216.76
Cd	0.00-1.89	0.00-6.67	0.04±0.09	0.17±0.36
Pb	0.00-1.70	0.00-7.10	0.20±0.29	0.78±1.11
Na	218.98 - 1246.23	927.78-4139.58	589.16 ± 247.20	2245.95 ± 837.72
K	294.93-1421.38	1382.35-5666.67	896.62 ± 268.85	3453.49 ± 1016.20
Ca	566.78-2780.44	2211.54-9927.50	1577.90 ± 569.04	6060.37 ± 2048.63
Mg	156.55-660.98	572.37-2360.00	307.12 ± 118.73	1172.86 ± 396.21

SD=Standard deviation

Table 2. Regression analysis of log wet body weight (g) versus log body burden element (µ/µg) for *Scomberoides commersonnianus*.

Elements	R	a	b	S.E (b)	t value (b = 1)
Fe	0.197 ^{ns}	64.840	-8.751	5.145	-8.946
Zn	0.522 ^{***}	248.692	-71.322	13.810	-71.394
Cd	0.044 ^{ns}	0.213	0.092	0.247	-3.957
Pb	0.283 [*]	-0.844	0.800	0.322	-14.961
Na	0.405 ^{***}	-94.727	257.565	68.993	257.550
K	0.262 [*]	414.942	181.407	79.197	181.394
Ca	0.386 ^{***}	75.860	565.694	160.209	565.688
Mg	0.287 [*]	74.256	87.701	34.717	87.672

*** = P<0.001, ** = P<0.01, * = P<0.05, ^{ns} = > 0.05. R =Regression coefficient, a=constant, b=Slope and SE=Standard error.

Table 3. Regression analysis of log total length (TL, cm) versus log body burden element (µ/µg) for *Scomberoides commersonnianus*.

Elements	R	a	b	S.E (b)	t value (b = 3)
Fe	0.206 ^{ns}	78.373	-23.180	13.065	-23.409
Zn	0.521 ^{***}	346.354	-180.947	35.158	-181.033
Cd	0.051 ^{ns}	0.033	0.268	0.628	-4.509
Pb	0.284 [*]	-1.962	2.044	0.819	-1.619
Na	0.391 ^{**}	-414.411	632.649	176.634	632.632
K	0.258 [*]	174.468	455.239	201.628	455.224
Ca	0.384 ^{***}	-690.161	1429.776	407.926	1429.768
Mg	0.280 [*]	-38.514	217.888	88.482	217.854

*** = P<0.001, ** = P<0.01, * = P<0.05 ^{ns} = > 0.05. =Regression coefficient, a=constant, b=Slope and SE=Standard error.

Table 4. Regression analysis of condition factor versus metal concentration (µgg⁻¹) in wet body weight for *Scomberoides commersonnianus*.

Elements	R	a	b	S.E (b)	t value (b=0)
Fe	0.628 ^{***}	83386.35	-73664.9	10832.82	-6.800
Zn	0.441 ^{***}	-61.951	151.420	36.514	4.146
Cd	0.057 ^{ns}	0.072	-0.035	0.072	-0.482
Pb	0.095 ^{ns}	0.345	-0.176	0.218	-0.805
Na	0.603 ^{***}	1483725	-1353911	212321.3	-6.376
K	0.711 ^{***}	2025078	-1819365	213106.2	-8.537
Ca	0.676 ^{***}	3572088	-3188308	412219.4	-7.734
Mg	0.609 ^{***}	709200.5	-636912	98322.75	-6.477

*** = P<0.001, ^{ns} = > 0.05. R =Regression coefficient, a=constant, b=Slope and SE=Standard error.

of condition factor and wet body weight with concentration of metal for *S. commersonnianus* are presented in Table 6. Concentration of K, Na, Mg, Ca and Fe showed highly significant positive correlation with total body length

Table 5. Multiple regression analysis of wet body weight (W, g.) and total length (TL, cm) with metal concentration (wet body weight, $\mu\text{g g}^{-1}$) for *Scomberoides commersonnianus*.

Relationship	R	a	$b_1 \pm \text{S.E}$	$b_2 \pm \text{S.E}$	VIF
$\text{Fe} = a + b_1W + b_2\text{TL}$	0.810***	16930.07	72.78 ± 19.07	-898.45 ± 546.72	0.343
$\text{Zn} = a + b_1W + b_2\text{TL}$	0.285*	-56173.8	-121.37 ± 56.29	3800.56 ± 1613.18	0.918
$\text{Cd} = a + b_1W + b_2\text{TL}$	0.456***	-0.286	-0.001 ± 0.0002	0.018 ± 0.004	0.792
$\text{Pb} = a + b_1W + b_2\text{TL}$	0.173 ^{ns}	-0.232	-0.001 ± 0.001	0.021 ± 0.014	0.970
$\text{Na} = a + b_1W + b_2\text{TL}$	0.872***	654762.6	2139.95 ± 303.948	-38067.2 ± 8710.11	0.237
$\text{K} = a + b_1W + b_2\text{TL}$	0.886***	190365.7	1408.11 ± 328.183	-11655.3 ± 9404.59	0.213
$\text{Ca} = a + b_1W + b_2\text{TL}$	0.827***	7286.755	1815.28 ± 735.31	-2150.58 ± 21071.54	0.315
$\text{Mg} = a + b_1W + b_2\text{TL}$	0.816***	182041.1	733.50 ± 167.38	-10480.7 ± 4796.78	0.332

a = Interceptb1 and b2 = Regression Coefficients for many factor to correlate among eachother. *** = $P < 0.001$. * = Least significant correlation. ^{ns} = > 0.05 .

Table 6. Multiple regression analysis of wet body weight (W, g.) and condition factor (K) with metal concentration (wet body weight, $\mu\text{g g}^{-1}$) for *Scomberoides commersonnianus*

Relationship	R	a	$b_1 \pm \text{S.E}$	$b_2 \pm \text{S.E}$	VIF
$\text{Fe} = a + b_1W + b_2K$	0.804***	-15099.7	47.087 ± 6.631	13713.16 ± 14852.47	0.352
$\text{Zn} = a + b_1W + b_2K$	0.160 ^{ns}	70434.85	-9.474 ± 19.893	-49342.1 ± 44557.87	0.974
$\text{Cd} = a + b_1W + b_2K$	0.335**	0.408	-0.0002 ± 0.0001	-0.333 ± 0.122	0.888
$\text{Pb} = a + b_1W + b_2K$	0.127 ^{ns}	0.606	-0.0001 ± 0.0002	-0.408 ± 0.391	0.984
$\text{Na} = a + b_1W + b_2K$	0.849***	-753816	1069.806 ± 112.738	631263.3 ± 252515.2	0.278
$\text{K} = a + b_1W + b_2K$	0.884***	-215720	1071.364 ± 113.479	168699.2 ± 254174.7	0.217
$\text{Ca} = a + b_1W + b_2K$	0.827***	-178019	1792.99 ± 252.212	138832.9 ± 564914.5	0.315
$\text{Mg} = a + b_1W + b_2K$	0.808***	-217757	443.194 ± 58.530	185496.1 ± 131097.6	0.345

a = Intercept; b_1 and b_2 = Regression Coefficients; SE = standard error; R = RegressionValue. *** = $P < 0.001$. ** = $P < 0.0$. ^{ns} = > 0.05 .

($P < 0.001$) in *S. commersonnianus* with multiple regression values ranging from 0.804 to 0.884; Cadmium showed least significant positive correlation; while Zn and Pb exhibited non-significant correlation ($P > 0.05$) with value.

4. Discussion

Accumulation of metals in fish was as in following order $\text{Ca} > \text{K} > \text{Na} > \text{Mg} > \text{Zn} > \text{Fe} > \text{Pb} > \text{Cd}$. Regression analysis of these metals were statistically analyzed with respect to condition factor and body size.

Metal concentrations in body of *S. commersonnianus* fish varied considerably. A minimum concentrations of heavy metal were found in Cd (wet and dry body weight) and Calcium in maximum concentration found the fish (wet and dry body weight) as the concentration of Calcium found maximum and cadmium in minimum concentration in *Oncorhynchus mykiss* (Naeem et al., 2010) and *Aristichthys nobilis* (Naeem et al., 2011). As this data on Calcium are also well-matched with study by Ansari et al. (2000) and Salam et al. (2002). Zhang et al. (2017) studied

that concentration of Pb, Zn and Cd as 0.011, 17.2 and 0.003 mg/kg, respectively in *Ctenopharyngodon idella* as almost similar values were reported in our study. Iron in present study were found agreement with previously reported studies (Canli&Atli 2003; Obasohan et al. 2006).

Permissible limit of Cadmium as recommended by the FAO (2011) as 0.05 as in present study within range. This implies Cd as not taken above as limit as by Ashraf and Nazeer (2010) and Chatta et al. (2016). Accumulations of Pb in study found below than maximum permissible limit as (FAO, 1984) reported similar concentration of lead in *Ctenopharyngodon idella* as in the present study. Zinc concentration in present study found within normal range by FAO (1989). The analyzed concentrations as noticed values agree with those reported by Shearer (1984) in *Oncorhynchus mykiss*, Salam and Naeem (1993) in *Tor putitora*, Salam and Davies (1994) in *Catla catla*, Naeem et al. (2011) in *Oreochromis niloticus*, Naeem et al. (2011) in *Mystus bleekeri* and Yousaf et al. (2012) in *Wallago attu*.

Sodium in dry and wet weight shown that concentration within normal range. Earlier, the mean concentration of

sodium documented in *Puntius sophore* 3335µg/g and *Trevor baim* 3746µg/g as (Rashid et al., 2012). Naeem et al. (2010) examined Potassium and described similar concentration in *O. mykiss* and *A. nobilis* (Naeem et al., 2011). The condition factor in this investigation remained consistent irrespective of the metal concentrations studied. Farkas et al. (2002) and Naeem et al. (2008) found similar results in other fish species. These findings demonstrate that the condition factor has no effect on the body size of the fish. According to the results of multiple regression analyses Cd, Zn, Fe, Na, Pb, K, Ca and Mg were shown to be substantially associated with total length, wet body weight and condition factor, as stated by (Naeem et al., 2011).

5. Conclusion

Conclusion of present study is that all the concentrations of various studied metals were found in permissible limit. Hence, the fish is safe for human consumption in these localities and it is likely to be an excellent source of vital metals. Moreover, condition factor of the fish body and fish size has a significant impact on metal accumulation. Findings of the study can be helpful for different organization for monitoring in future.

References

- ABBAS, S., IQBAL, A., ANJUM, K.M., SHERZADA, S., ATIQUE, U., KHAN, M.K.A., AKMAL, M., RAHMAN, A., ASIF, A.R., AHMAD, S., MALIK, A., KHAN, S.A., AHMAD, S. and INAYAT, M., 2021. Body composition, growth performance and enzyme activities of *Labeo rohita* fed different commercial fish feeds. *Brazilian Journal of Biology = Revista Brasileira de Biologia*, vol. 83, pp. e250402. <http://dx.doi.org/10.1590/1519-6984.250402>. PMID:34614125.
- AHMAD, B., HUSSAIN, S.M., ALI, S., ARSALAN, M., TABASSUM, S. and SHARIF, A., 2021. Efficacy of acidified phytase supplemented cottonseed meal based diets on growth performance and proximate composition of *Labeo rohita* fingerlings. *Brazilian Journal of Biology = Revista Brasileira de Biologia*, vol. 83, pp. e247791. <http://dx.doi.org/10.1590/1519-6984.247791>. PMID:34431918.
- ANSARI, T.M., SALAM, A. and ASIF, M., 2000. Inorganic elements in relation to growth in farmed silver carp, (*Hypophthalmichthys molitrix*). *Pakistan Journal of Fisheries*, vol. 1, pp. 81.
- ASEGBOLOYI, J.N., ONYIMONYI, A.E., UJAM, O.T., UKWUEZE, N.N. and UKOHA, P.O., 2010. Assessment of toxic trace metals in selected fish species and parts of domestic animals. *Pakistan Journal of Nutrition*, vol. 9, no. 3, pp. 213-215. <http://dx.doi.org/10.3923/pjn.2010.213.215>.
- ASHRAF, W. and NAZEER, Z., 2010. Heavy metals burden in kidney and heart tissues of *Scarus ghobban* fish from the Eastern province of Saudi Arabia. *Bulletin of the Chemical Society of Ethiopia*, vol. 24, no. 1, pp. 139-143. <http://dx.doi.org/10.4314/bcse.v24i1.52977>.
- BARSE, A.V., CHAKRABARTI, T., GHOSH, T.K., PAL, A.K. and JADHAO, S.B., 2007. Endocrine disruption and metabolic changes following exposure of *Cyprinus carpio* to diethyl phthalate. *Pesticide Biochemistry and Physiology*, vol. 88, no. 1, pp. 36-42. <http://dx.doi.org/10.1016/j.pestbp.2006.08.009>.
- CANLI, M. and ATLI, G., 2003. The relationships between heavy metal (Cd, Cr, Cu, Fe, Pb, Zn) levels and the size of six Mediterranean fish species. *Environmental Pollution*, vol. 121, no. 1, pp. 129-136. [http://dx.doi.org/10.1016/S0269-7491\(02\)00194-X](http://dx.doi.org/10.1016/S0269-7491(02)00194-X). PMID:12475070.
- CHATTA, A., KHAN, M., MIRZA, Z. and ALI, A., 2016. Heavy metal (cadmium, lead, and chromium) contamination in farmed fish: a potential risk for consumers' health. *Turkish Journal of Zoology*, vol. 40, pp. 248-256. <http://dx.doi.org/10.3906/zoo-1506-1>.
- COETZEE, L., 1998. *Bioaccumulation of metals in selected fish species and the effects of pH on aluminum toxicity in a cichlid O. mossambicus*. Johannesburg, South Africa: Rand Afrikaans University. PhD Dissertation.
- EISLER, R., 1987. *Mercury hazards to fish, wildlife and invertebrates: a synoptic review*. Washington, D.C.: Fish Wild Life Services.
- EVERAERT, G., VAN CAUWENBERGHE, L., DE RIJCKE, M., KOELMANS, A.A., MEES, J., VANDEGEHUCHTE, M. and JANSSEN, C.R., 2018. Risk assessment of microplastics in the ocean: modelling approach and first conclusions. *Environmental Pollution*, vol. 242, no. Pt B, pp. 1930-1938. <http://dx.doi.org/10.1016/j.envpol.2018.07.069>. PMID:30061084.
- FOOD AND AGRICULTURE ORGANIZATION – FAO, and WORLD HEALTH ORGANIZATION – WHO. *Evaluation of certain food additives and the contaminants mercury, lead and cadmium*. Rome: FAO, 1989. WHO Technical Report, no. 505.
- FARKAS, A., SALANKI, J. and SPECZIÁR, A., 2002. Relation between growth and the heavy metal concentration in organs of bream *Abramis brama* L. Populating Lake Balaton. *Archives of Environmental Contamination and Toxicology*, vol. 43, no. 2, pp. 236-243. <http://dx.doi.org/10.1007/s00244-002-1123-5>. PMID:12115050.
- GASPERI, J., WRIGHT, S.L., DRIS, R., COLLARD, F., MANDIN, C., GUERROUACHE, M., LANGLOIS, V., KELLY, F.J. and TASSIN, B., 2018. Microplastics in air: are we breathing it in? *Current Opinion in Environmental Science Health*, vol. 1, pp. 1-5. <http://dx.doi.org/10.1016/j.coesh.2017.10.002>.
- KOELMANS, A.A., KOOI, M., LAW, K.L. and VAN SEBILLE, E., 2017. All is not lost: deriving a top-down mass budget of plastic at sea. *Environmental Research Letters*, vol. 12, no. 11, pp. 114028. <http://dx.doi.org/10.1088/1748-9326/aa9500>.
- NAEEM, M., HASANAIN, S.K. and MUMTAZ, A., 2008. Electrical transport and optical studies of ferromagnetic cobalt doped ZnO nanoparticles exhibiting a metal-insulator transition. *Journal of Physics Condensed Matter*, vol. 20, no. 2, pp. 025210. <http://dx.doi.org/10.1088/0953-8984/20/2/025210>.
- NAEEM, M., SALAM, A., TAHIR, S.S. and RAUF, N., 2010. Assessment of the essential and toxic heavy metals in hatchery reared *Oncorhynchus mykiss*. *International Journal of Agriculture and Biology*, vol. 12, pp. 935-938.
- NAEEM, M., SALAM, A., TAHIR, S.S. and RAUF, N., 2011. Effect of fish size and condition on the contents of twelve essential and non-essential elements in *Aristichthys nobilis*. *Pakistan Veterinary Journal*, vol. 31, pp. 109-112.
- NASIR, N.A. and AL-NAJAR, G.A., 2015. Seasonal variations of heavy metal concentrations in tissues of talang queenfish (*Scomberoides commersonianus*) from Iraqi Marine and Coastal Waters, Northwestern Arabian Gulf. *Mesopotamian Journal of Marine Science*, vol. 30, pp. 112-123.
- OBASOHAN, E.E., ORONSAYE, J.A.O. and OBANO, E.E., 2006. Heavy metal concentrations in *Malapterurus electricus* and *Chrysichthys nigrodigitatus* from Ogba River in Benin City, Nigeria. *African Journal of Biotechnology*, vol. 5, pp. 974-982.
- RAHMAN, M.S., MOLLA, A.H., SAHA, N. and RAHMAN, A., 2012. Study on heavy metals levels and its risk assessment in some edible fishes from Bangshi River, Dhaka, Bangladesh. *Food Chemistry*,

- vol. 134, no. 4, pp. 1847-1854. <http://dx.doi.org/10.1016/j.foodchem.2012.03.099>. PMID:23442629.
- RASHID, H., HASAN, M.N., TANU, M.B., PARVEEN, R., SUKHAN, Z.P., RAHMAN, M.S. and MAHMUD, Y., 2012. Heavy metal pollution and chemical profile of Khiru River, Bangladesh. *International Journal of Environment*, vol. 2, pp. 57-63.
- SALAM, A. and DAVIES, P.M.C., 1994. Body composition of northern pike (*Esox lucius* L.) in relation to body size and condition factor. *Fisheries Research*, vol. 19, no. 3-4, pp. 193-204. [http://dx.doi.org/10.1016/0165-7836\(94\)90038-8](http://dx.doi.org/10.1016/0165-7836(94)90038-8).
- SALAM, A., ANSARI, T.M., TARIC, N. and AKHTAR, Q.A., 2002. Effect of body size on metal concentrations in farmed 334.-*Cirrhinus mrigala*. *Asian Fisheries Science*, vol. 15, no. 4, pp. 329-334. <http://dx.doi.org/10.33997/j.afs.2002.15.4.004>.
- SALAM, A.M. and NAEEM, Z.B., 1993. Length-weight and condition factor relationship of fresh water Chinese carp, from fish hatchery Islamabad. *Pakistan Journal of Research*, vol. 5, pp. 45-48.
- SHEARER, K.D., 1984. Changes in the elemental composition of hatchery reared rainbow trout (*Salmo gairdneri*) associated with growth and reproduction. *Canadian Journal of Fisheries and Aquatic Science*, vol. 41, no. 11, pp. 1592-1600. <http://dx.doi.org/10.1139/f84-197>.
- STANISKIENE, B., MATUSEVICIUS, P. and URBONAVIČIUS, A., 2009. Distribution of heavy metals in muscles of fish: concentrations and change tendencies. *Environmental Research, Engineering and Management*, vol. 48, pp. 35-41.
- THE FOOD AND AGRICULTURE ORGANIZATION – FAO, 2011. *FAO Fisheries and Aquaculture Report No. 978*. Rome: FAO.
- THE FOOD AND AGRICULTURE ORGANIZATION – FAO, 1984. *List of maximum levels recommended for contaminants by the Joint. Rome/Geneva*: FAO/WHO.
- YOUSAF, M., NAEEM, M. and KHOKHAR, M.Y., 2012. Effect of body size on elemental concentration in wild *Wallago attu* (Block and Schneider) from southern Punjab, Pakistan. *African Journal of Biotechnology*, vol. 11, pp. 1764-1767.
- ZHANG, J., ZHU, L., LI, F., LIU, C., YANG, Z., QIU, Z. and XIAO, M., 2017. Heavy metals and metalloid distribution in different organs and health risk assessment for edible tissues of fish captured from Honghu Lake. *Oncotarget*, vol. 8, no. 60, pp. 101672-101685. <http://dx.doi.org/10.18632/oncotarget.21901>. PMID:29254195.